

Spectral solutions to stochastic models of gene expression with bursts and regulation

Andrew Mugler*

Department of Physics, Columbia University, New York, NY 10027

Aleksandra M. Walczak†

Princeton Center for Theoretical Science, Princeton University, Princeton, NJ 08544

Chris H. Wiggins‡

Department of Applied Physics and Applied Mathematics,

Center for Computational Biology and Bioinformatics, Columbia University, New York, NY 10027

(Dated: July 20, 2009)

Signal-processing molecules inside cells are often present at low copy number, which necessitates probabilistic models to account for intrinsic noise. Probability distributions have traditionally been found using simulation-based approaches which then require estimating the distributions from many samples. Here we present in detail an alternative method for directly calculating a probability distribution by expanding in the natural eigenfunctions of the governing equation, which is linear. We apply the resulting *spectral* method to three general models of stochastic gene expression: a single gene with multiple expression states (often used as a model of bursting in the limit of two states), a gene regulatory cascade, and a combined model of bursting and regulation. In all cases we find either analytic results or numerical prescriptions that greatly outperform simulations in efficiency and accuracy. In the last case, we show that bimodal response in the limit of slow switching is not only possible but optimal in terms of information transmission.

Signals are processed in cells using networks of interacting components, including proteins, mRNAs, and small signaling molecules. These components are usually present in low numbers [1, 2, 3, 4, 5, 6], which means the size of the fluctuations in their copy counts is comparable to the copy counts themselves. Noise in gene networks has been shown to propagate [7], and therefore explicitly accounting for the stochastic nature of gene expression appears important when predicting the properties of real biological networks.

Although summary statistics such as mean and variance are sometimes sufficient for answering questions of biological interest [8], calculating certain quantities, such as information transmission [9, 10, 11, 12, 13, 14] or escape properties [15, 16, 17, 18, 19], requires knowing the full probability distribution. Full knowledge of the probability distribution can also be used to discern different molecular models of the noise sources based on recent exact measurements of probability distributions [20, 21, 22, 23].

Much analytical and purely computational effort has gone into detailed models of noise in small genetic switches [8, 24, 25, 26]. The most general description is based on the master equation describing the time evolution of the joint probability distribution over all copy counts [27]. Some progress has been made by applying approximations to the master equation. For example, a wide class of approximations focuses on limits of large

concentrations or small switches [8, 24, 28]. Approximations based on timescale separation of the steps of small signaling cascades have also been successfully used to calculate escape properties [15, 29, 30]. More often, modelers resort to stochastic simulation techniques, the most common of which is based on the varying-step Monte Carlo method [31, 32]. Probabilistic modeling of stochastic systems by simulation requires a computational challenge (generating many sample trajectories) followed by an even more difficult statistical challenge (parameterizing or otherwise estimating the probability distribution from which the samples are drawn) [33].

In a recent paper [34] we introduce a new method for calculating the steady state distributions of chemical reactants, which we call the spectral method. The procedure relies on exploiting the natural basis of a simpler problem from the same class. The full problem is then solved numerically as an expansion in the natural basis [48]. In the spectral method we use the analytical guidance of a simple birth-death problem to reduce the master equation for a cascade to a set of algebraic equations. We break the problem into two parts: a parameter-independent preprocessing step, and the parameter-dependent step of obtaining the actual probability distributions. The spectral method allows huge computational gains with respect to simulations. In prior work [34] we illustrate the method in the example of gene regulatory cascades. We combine the spectral method with a Markov approximation, which exploits the observation that the behavior of a given species should depend only weakly on distant nodes given the proximal nodes.

In this paper we expand upon the application of the spectral method to more biologically realistic models of regulation: (i) a model of bursting in gene expression

*ajm2121@columbia.edu

†awalczak@princeton.edu

‡chris.wiggins@columbia.edu

and (ii) a model that includes both bursts and explicit regulation by binding of transcription factor proteins. In both cases we demonstrate how the spectral method gives either analytic results or reduced algebraic expressions that can be solved numerically in orders of magnitude less time than stochastic simulations.

We begin with a model of a multi-state birth-death process, a special case of which has been used to describe transcriptional bursting [22, 35]. We illustrate how the spectral method reduces the model to a simple iterative algebraic equation, and in the appropriate limiting case recovers the known analytic results. We also use this section to introduce the basic notation used throughout the paper. Next we explore the problem of gene regulation in detail. The main idea behind the spectral method is the exploitation of an underlying natural basis for a problem which we can solve exactly. We explore four different spectral representations of the regulation model used in previous work [34] that arise from four natural choices of eigenbasis in which to expand the solution (cf. Sec. II A and Fig. 2). All representations reduce the master equation to a set of linear algebraic equations, and one admits an analytic solution by virtue of the tridiagonal matrix algorithm. We compare the efficiencies of the representations' numerical implementations and show that all outperform simulation. Lastly, we apply the spectral method to a model that combines bursting and regulation. We obtain a linear algebraic expression that permits large speedup over simulation and thus admits optimization of information transmission. Optimization reveals two types of solutions: a unimodal response when the rates of switching between expression states are comparable to degradation rates, and a bimodal response when switching rates are much slower than degradation rates.

I. BURSTS OF GENE EXPRESSION

We first consider a model of gene expression in which a gene exists in one of Z stochastic “states,” i.e. protein production obeys a simple birth-death process, but with a state-dependent birth rate. In the special case of $Z = 2$, this corresponds to a gene existing in an on- or an off-state due, for example, to the binding and unbinding of the RNA polymerase. Such a model has been used to describe transcriptional bursting [22, 35], and we specialize to this case in Sec. I C.

For a general Z -state process, the master equation

$$\dot{p}_n^z = g_z p_{n-1}^z + (n+1)p_{n+1}^z - (g_z + n)p_n^z + \sum_{z'} \Omega_{zz'} p_n^{z'} \quad (1)$$

describes the time evolution of the joint probability distribution p_n^z , where z specifies the state ($1 \leq z \leq Z$), n is the number of proteins, g_z is the production rate in state z , $\Omega_{zz'}$ is a stochastic matrix of transition rates between states, and $\dot{p}_n^z$ denotes differentiation of the probability distribution with respect to time. Time and all rates

have been nondimensionalized by the protein degradation rate. Note that conservation of probability requires

$$\sum_z \Omega_{zz'} = 0. \quad (2)$$

The relationship between the transition rates in $\Omega_{zz'}$ and the probabilities $\pi_z = \sum_n p_n^z$ of being in the z th state can be seen by summing Eqn. 1 over n ; at steady state one obtains

$$\sum_{z'} \Omega_{zz'} \pi_{z'} = 0, \quad (3)$$

and normalization requires

$$\sum_z \pi_z = 1. \quad (4)$$

In the following sections, we introduce the spectral method and demonstrate how it can be used to solve for the full joint distribution p_n^z .

A. Notation and definitions

We begin our solution of Eqn. 1 by defining the generating function [27] $G_z(x) = \sum_n p_n^z x^n$ over complex variable x [49] (note that superscript z is an index, while superscript n on x^n is a power). It will prove more convenient to rewrite the generating function in a more abstract representation using states indexed by protein number $|n\rangle$,

$$|G_z\rangle = \sum_n p_n^z |n\rangle, \quad (5)$$

with inverse transform

$$p_n^z = \langle n | G_z \rangle. \quad (6)$$

The more familiar form can be recovered by projecting the position space $\langle x |$ onto Eqn. 5, with the provision that

$$\langle x | n \rangle = x^n. \quad (7)$$

Concurrent choices of conjugate state

$$\langle n | x \rangle = \frac{1}{x^{n+1}} \quad (8)$$

and inner product

$$\langle f | f' \rangle = \oint \frac{dx}{2\pi i} \langle f | x \rangle \langle x | f' \rangle \quad (9)$$

ensure orthonormality of the states, $\langle n | n' \rangle = \delta_{nn'}$, as can be verified using Cauchy's theorem,

$$\oint \frac{dx}{2\pi i} \frac{f(x)}{(x - x_0)^{n+1}} = \frac{1}{n!} \partial_x^n [f(x)]_{x=x_0} \theta(n+1), \quad (10)$$

where the convention $\theta(0) = 0$ is used for the Heaviside function.

With these definitions, summing Eqn. 1 over n against $|n\rangle$ yields

$$|\dot{G}_z\rangle = -(\hat{a}^+ - 1)(\hat{a}^- - g_z)|G_z\rangle + \sum_{z'} \Omega_{zz'}|G_{z'}\rangle, \quad (11)$$

where the operators $\hat{a}^+$ and $\hat{a}^-$ raise and lower protein number, respectively, i.e.

$$\hat{a}^+|n\rangle = |n+1\rangle, \quad (12)$$

$$\hat{a}^-|n\rangle = n|n-1\rangle, \quad (13)$$

with adjoint operations

$$\langle n|\hat{a}^+ = \langle n-1|, \quad (14)$$

$$\langle n|\hat{a}^- = (n+1)\langle n+1|. \quad (15)$$

As in the operator treatment of the simple harmonic oscillator in quantum mechanics, the raising and lower operators satisfy the commutation relation $[\hat{a}^-, \hat{a}^+] = 1$, and $\hat{a}^+\hat{a}^-$ is a number operator, i.e. $\hat{a}^+\hat{a}^-|n\rangle = n|n\rangle$ [50]. This operator formalism for the generating function was introduced in the context of diffusion independently by Doi [36] and Zeldovich [37], and developed by Peliti [38]. A review by Mattis and Glasser [39] introduces and discusses the applications of the formalism for diffusion.

The factorized form of the birth-death operator in Eqn. 11 suggests the definition of shifted raising and lowering operators

$$\hat{b}^+ = \hat{a}^+ - 1 \quad (16)$$

$$\hat{b}_z^- = \hat{a}^- - g_z, \quad (17)$$

making Eqn. 11

$$|\dot{G}_z\rangle = -\hat{b}^+\hat{b}_z^-|G_z\rangle + \sum_{z'} \Omega_{zz'}|G_{z'}\rangle. \quad (18)$$

Since $\hat{b}^+\hat{b}_z^-$ is a new number operator, it is clear that the eigenvalues of the birth-death operator $-\hat{b}^+\hat{b}_z^-$ are nonpositive integers, i.e.

$$\hat{b}^+\hat{b}_z^-|j_z\rangle = j|j_z\rangle, \quad (19)$$

where nonnegative integers j index (z -dependent) eigenfunctions $|j_z\rangle$. In position space the j th eigenfunction is

$$\langle x|j_z\rangle = (x-1)^j e^{g_z(x-1)}, \quad (20)$$

with conjugate

$$\langle j_z|x\rangle = \frac{e^{-g_z(x-1)}}{(x-1)^{j+1}}, \quad (21)$$

such that orthonormality $\langle j_z|j'_z\rangle = \delta_{jj'}$ is satisfied under the inner product in Eqn. 9. Note that $\hat{b}^+$ and $\hat{b}_z^-$ raise

and lower eigenstates $|j_z\rangle$ as in Eqns. 12-15, i.e.

$$\hat{b}^+|j_z\rangle = |(j+1)_z\rangle, \quad (22)$$

$$\hat{b}_z^-|j_z\rangle = j|(j-1)_z\rangle, \quad (23)$$

$$\langle j_z|\hat{b}^+ = \langle (j-1)_z|, \quad (24)$$

$$\langle j_z|\hat{b}_z^- = (j+1)\langle (j+1)_z|. \quad (25)$$

As we will see in this and subsequent sections, going between the protein number basis $|n\rangle$ and the eigenbasis $|j_z\rangle$ requires the mixed product $\langle n|j_z\rangle$ or its conjugate $\langle j_z|n\rangle$. There are several ways of computing these objects, as described in Appendix A. Notable special cases are

$$\langle (j=0)_z|n\rangle = 1, \quad (26)$$

$$\langle n|(j=0)_z\rangle = e^{-g_z} \frac{(g_z)^n}{n!}, \quad (27)$$

where the latter is the Poisson distribution.

B. The spectral method

We now demonstrate how the spectral method exploits the eigenfunctions $|j_z\rangle$ to decompose and simplify the equation of motion. Expanding the generating function in the eigenbasis,

$$|G_z\rangle = \sum_j G_j^z |j_z\rangle, \quad (28)$$

and projecting the conjugate state $\langle j_z|$ onto Eqn. 18 yields the equation of motion

$$\dot{G}_j^z = -jG_j^z + \sum_{z'} \Omega_{zz'} \sum_{j'} G_{j'}^{z'} \langle j_z|j'_{z'}\rangle \quad (29)$$

for the expansion coefficients G_j^z (where the dummy index j in Eqn. 28 has been changed to j' in Eqn. 29). Using Eqns. 9, 10, 20, and 21, the product $\langle j_z|j'_{z'}\rangle$ evaluates to

$$\langle j_z|j'_{z'}\rangle = \frac{(-\Delta_{zz'})^{j-j'}}{(j-j')!} \theta(j-j'+1), \quad (30)$$

where $\Delta_{zz'} = g_z - g_{z'}$. Noting that $\langle j_z|j_{z'}\rangle = 1$ and that $\langle j_z|j'_{z'}\rangle = 0$ for $j' < j$, Eqn. 29 becomes

$$\begin{aligned} \dot{G}_j^z &= -jG_j^z + \sum_{z'} \Omega_{zz'} G_{j'}^{z'} \\ &+ \sum_{z' \neq z} \Omega_{zz'} \sum_{j' < j} G_{j'}^{z'} \frac{(-\Delta_{zz'})^{j-j'}}{(j-j')!}. \end{aligned} \quad (31)$$

The last term in Eqn. 31 makes clear that each j th term is slaved to terms with $j' < j$, allowing the G_j^z to be computed iteratively in j . The lower-triangular structure of the equation is a consequence of rotating to the

eigenspace of the birth-death operator; this structure was not present in the original master equation.

At steady-state, G_j^z obeys

$$jG_j^z - \sum_{z'} \Omega_{zz'} G_j^{z'} = \sum_{z' \neq z} \Omega_{zz'} \sum_{j' < j} G_{j'}^{z'} \frac{(-\Delta_{zz'})^{j-j'}}{(j-j')!}, \quad (32)$$

from which it is clear that G_j^z can be computed successively in j . Since

$$G_0^z = \langle (j=0)_z | G_z \rangle = \sum_n p_n^z \langle (j=0)_z | n \rangle = \pi_z \quad (33)$$

(cf. Eqn. 26), the computation is initialized using Eqns. 3-4, i.e.

$$\sum_{z'} \Omega_{zz'} G_0^{z'} = 0, \quad (34)$$

$$\sum_z G_0^z = 1. \quad (35)$$

Recalling Eqns. 6 and 28, the probability distribution is retrieved via

$$p_n^z = \sum_j G_j^z \langle n | j_z \rangle, \quad (36)$$

where the mixed product $\langle n | j_z \rangle$ can be computed as described in Appendix A.

There is an alternative way to decompose the master equation spectrally. Instead of expanding the generating function in eigenfunctions $|j_z\rangle$, which depend on the production rates g_z in each state, we may expand in eigenfunctions parameterized by a single rate $\bar{g}$, i.e.

$$|G_z\rangle = \sum_j G_j^z |j\rangle, \quad (37)$$

where

$$\langle x | j \rangle = (x-1)^j e^{\bar{g}(x-1)}, \quad (38)$$

$$\langle j | x \rangle = \frac{e^{-\bar{g}(x-1)}}{(x-1)^{j+1}}. \quad (39)$$

The parameter $\bar{g}$ is arbitrary and thus acts as a “gauge” freedom.

We may now partition the birth-death operator as

$$-\hat{b}^+ \hat{b}_z^- = -\hat{b}^+ \bar{b}^- - \hat{b}^+ \Gamma_z \quad (40)$$

where $\bar{b}^- = \hat{a}^- - \bar{g}$ such that the $|j\rangle$ are the eigenstates of $\hat{b}^+ \bar{b}^-$, i.e.

$$\hat{b}^+ \bar{b}^- |j\rangle = j |j\rangle, \quad (41)$$

and $\Gamma_z = \bar{g} - g_z$ describes the deviation of each state’s production rate from the constant $\bar{g}$.

Projecting the conjugate state $\langle j|$ onto Eqn. 18 and using Eqn. 37 (with dummy index j changed to j') gives

$$\dot{G}_j^z = -jG_j^z + \sum_{j'} \langle j | \hat{b}^+ \Gamma_z | j' \rangle G_{j'}^z + \sum_{z'} \Omega_{zz'} G_j^{z'}. \quad (42)$$

Recalling Eqn. 24, Eqn. 42 at steady state becomes

$$\sum_{z'} (\Omega_{zz'} - j\delta_{zz'}) G_j^{z'} = \Gamma_z G_{j-1}^z. \quad (43)$$

Eqn. 43 is subdiagonal in j , meaning computation of the j th term requires only the previous $(j-1)$ th term and the inversion of the Z -by- Z matrix $(\mathbf{\Omega} - j\mathbf{I})$ (where $\mathbf{I}$ is the identity matrix). It is initialized with Eqns. 34-35 and solved successively in j . The probability distribution is retrieved via

$$p_n^z = \sum_j G_j^z \langle n | j \rangle, \quad (44)$$

where $\langle n | j \rangle$ is computed as described in Appendix A.

As an example of a simple computation employing the spectral method, Fig. 1 shows probability distributions for the case of $Z = 3$ states, corresponding to a gene that is either off, producing proteins at a low rate, or producing proteins at a high rate. For simplicity we set the rates of switching among all states equal to a constant ω , making the stochastic matrix

$$\mathbf{\Omega} = \begin{pmatrix} -2\omega & \omega & \omega \\ \omega & -2\omega & \omega \\ \omega & \omega & -2\omega \end{pmatrix}. \quad (45)$$

As seen in Fig. 1, when $\omega \ll 1$ (corresponding to a switching rate much slower than the degradation rate) the dwell time in each expression state lengthens. The slow switching gives the protein copy number time to equilibrate in any of the three expression states, resulting in a trimodal marginal distribution p_n . When $\omega \gg 1$ (corresponding to a switching rate much faster than the degradation rate), the system switches frequently among the three expression states, resulting in an average production rate. In this limit, the expression state equilibrates on a faster timescale than the protein number state.

C. The on/off gene

For the special case of $Z = 2$ states, as when a gene is either “on” ($z = +$) or “off” ($z = -$), it is useful to demonstrate how the spectral method reproduces known analytic results [22, 35]. The probability distribution can be written in vector form as

$$\vec{p}_n = \begin{pmatrix} p_n^- \\ p_n^+ \end{pmatrix}, \quad (46)$$

and defining ω_+ and ω_- as the transition rates to and from the on-state, respectively, the stochastic matrix takes the form

$$\mathbf{\Omega} = \begin{pmatrix} -\omega_+ & \omega_- \\ \omega_+ & -\omega_- \end{pmatrix}. \quad (47)$$

Note that Eqn. 3 implies

$$\frac{\pi_-}{\pi_+} = \frac{\omega_-}{\omega_+}, \quad (48)$$




FIG. 1: An example of the spectral method for $Z = 3$ states. Dotted curves show the joint distribution p_n^z for each of $z = 1, 2$, and 3 , and solid curves show the marginal distribution p_n . The stochastic transition matrix is given in Eqn. 45, and the setting of ω in each panel is indicated in the upper-right corner. Production rates are $g_1 = 0$, $g_2 = 3$, and $g_3 = 12$ for all panels. Distributions are calculated via the spectral decomposition in Eqn. 43 with $\bar{g} = \langle g_z \rangle = 5$.

which makes clear that increasing the rate of transition to either state increases the probability of being in that state.

From Eqn. 32 the spectral expansion coefficients obey

$$jG_j^\pm + \omega_\mp G_j^\pm - \omega_\pm G_j^\mp = \omega_\pm \sum_{j' < j} G_{j'}^\mp \frac{(\mp\Delta)^{j-j'}}{(j-j')!}, \quad (49)$$

where $\Delta = \Delta_{+-} = -\Delta_{-+}$. Initializing with $G_0^\pm = \omega_\pm/(\omega_+ + \omega_-)$ and computing the first few terms reveals the pattern

$$G_j^\pm = \frac{\omega_\pm}{\omega_+ + \omega_-} \frac{(\mp\Delta)^j}{j!} \frac{\prod_{j'=0}^{j-1} (j' + \omega_\mp)}{\prod_{j''=0}^{j-1} (j'' + \omega_+ + \omega_- + 1)} \quad (50)$$

$$= \frac{\omega_\pm}{\omega_+ + \omega_-} \frac{(\mp\Delta)^j}{j!} \frac{\Gamma(j + \omega_\mp)}{\Gamma(\omega_\mp)} \frac{\Gamma(\omega_+ + \omega_- + 1)}{\Gamma(j + \omega_+ + \omega_- + 1)}, \quad (51)$$

where in the second line the products are written in terms of the Gamma function.

For comparison with known results [22, 35] we write the total generating function $|G\rangle = \sum_\pm |G_\pm\rangle$ in position space,

$$G(x) = \sum_\pm \langle x | G_\pm \rangle = \sum_\pm \sum_j \langle x | j_\pm \rangle \langle j_\pm | G_\pm \rangle \quad (52)$$

$$= \sum_\pm \sum_j (x-1)^j e^{g_\pm(x-1)} G_j^\pm \quad (53)$$

$$= \sum_\pm \frac{\omega_\pm}{\omega_+ + \omega_-} e^{g_\pm(x-1)} \times \Phi[\omega_\mp, \omega_+ + \omega_- + 1; \mp\Delta(x-1)], \quad (54)$$

where

$$\Phi[\alpha, \beta; y] = \sum_j \frac{\Gamma(j + \alpha)}{\Gamma(\alpha)} \frac{\Gamma(\beta)}{\Gamma(j + \beta)} \frac{y^j}{j!} \quad (55)$$

is the confluent hypergeometric function. As shown in Appendix B, in the limit $g_- = 0$, Eqn. 54 reduces to

$$G(x) = \Phi[\omega_+, \omega_+ + \omega_-; g_+(x-1)], \quad (56)$$

and the marginal p_n is given by

$$p_n = \frac{g_+^n}{n!} \frac{\Gamma(n + \omega_+)}{\Gamma(\omega_+)} \frac{\Gamma(\omega_+ + \omega_-)}{\Gamma(n + \omega_+ + \omega_-)} \times \Phi[\omega_+ + n, \omega_+ + \omega_- + n; -g_+], \quad (57)$$

in agreement with the results of Iyer-Biswas et al. [35] and Raj et al. [22]. We remind the reader that in addition to reducing to known results in the special case of $Z = 2$ states with a vanishing off-rate, the spectral method is valid for any number of states with arbitrary production rates.

II. GENE REGULATION

Next we consider a two-gene regulatory cascade, in which the production rate of the second gene is a function of the number of proteins of the first gene. As shown in previous work [34], a cascade of any length can be reduced to such a generalized two-dimensional system using the Markov approximation, which asserts that the probability distribution for a given node of the cascade should depend only weakly on the probability distributions of the distant modes given the proximal nodes.

In the present section, we consider only the generalized two-dimensional equation and explore different approaches to solving it. The equation describes two genes, each with one expression state, with regulation encoded by a functional dependence of the downstream protein production rate on the upstream protein copy number. In section IID we make an explicit connection between the on/off gene discussed in section IC and the case when the functional dependence is a threshold. Finally, in section III we combine the two types of models and consider a system with regulation and bursts.

A. Representations of the master equation

1. The $|n, m\rangle$ basis

Defining n and m as the numbers of proteins produced by the first and second gene, respectively, the master equation describing the time evolution of the joint probability distribution p_{nm} is [34]

$$\dot{p}_{nm} = g_{n-1}p_{n-1,m} + (n+1)p_{n+1,m} - (g_n + n)p_{nm} + \rho[q_n p_{n,m-1} + (m+1)p_{n,m+1} - (q_n + m)p_{nm}]. \quad (58)$$




FIG. 2: Summary of the bases discussed in Sec. IIA (black) and their gauge freedoms (barred parameters in grey). In the top row, neither the m nor the n sector is expanded in an eigenbasis; in the middle row, one sector is expanded; and in the bottom row, both sectors are expanded. The $|j, k\rangle$ basis can be viewed as a special case of the $|j, k_j\rangle$ or $|j, k_n\rangle$ basis with $\bar{q}_j = \bar{q}$ or $\bar{q}_n = \bar{q}$ respectively. The $|j, m\rangle$ basis is not discussed as it is not useful in simplifying the problem.

The function q_n describes the regulation of the second species by the first, and the function g_n describes the effective autoregulation of the first species, due either to a non-Poissonian input distribution or to effects further upstream in the case of a longer cascade [34]. Time is rescaled by the first gene's degradation rate, so that each gene's production rate (g_n or q_n) is normalized by its respective degradation rate, and ρ is ratio of the second gene's degradation rate to that of the first. We impose no constraints on the form of g_n or q_n —they can be arbitrary nonlinear functions.

Summing Eqn. 58 over m gives a simple recursion relation between g_n and p_n at steady state, from which explicit relations are easily identified. If p_n is known, g_n is found as

$$g_n = (n+1) \frac{p_{n+1}}{p_n}. \quad (59)$$

If on the other hand g_n is known, p_n is found as

$$p_n = \frac{p_0}{n!} \prod_{n'=0}^{n-1} g_{n'}, \quad (60)$$

with p_0 set by normalization. Note that if the first species obeys a simple birth-death process, $g_n = g = \text{constant}$, and Eqn. 60 reduces to the Poisson distribution.

In the current representation (Eqn. 58), which we denote the $|n, m\rangle$ basis, finding the steady state solution for the joint distribution p_{nm} means finding the null space of an infinite (or, effectively for numerical purposes, very large) locally banded tridiagonal matrix. More precisely, defining N as the numerical cutoff in protein number n or m , the problem amounts to inverting an N^2 -by- N^2 matrix, which is computationally taxing even for moderate cutoffs N .

In order to solve Eqn 58 more efficiently we will employ the spectral method. We begin as before by defining the

generating function [27] $G(x, y) = \sum_{nm} p_{nm} x^n y^m$ over complex variables x and y , or, in state notation,

$$|G\rangle = \sum_{nm} p_{nm} |n, m\rangle, \quad (61)$$

with inverse transform

$$p_{nm} = \langle n, m | G \rangle. \quad (62)$$

Summing Eqn. 58 over n and m against $|n, m\rangle$ and employing the same operator notation as in Eqns. 12-15 yields

$$|\dot{G}\rangle = -\hat{H}|G\rangle, \quad (63)$$

where

$$\hat{H} = \hat{b}_n^+ \hat{b}_n^-(n) + \rho \hat{b}_m^+ \hat{b}_m^-(n), \quad (64)$$

and

$$\hat{b}_n^+ = \hat{a}_n^+ - 1, \quad (65)$$

$$\hat{b}_m^+ = \hat{a}_m^+ - 1, \quad (66)$$

$$\hat{b}_n^-(n) = \hat{a}_n^- - \hat{g}_n, \quad (67)$$

$$\hat{b}_m^-(n) = \hat{a}_m^- - \hat{q}_n. \quad (68)$$

Here the regulation functions have been promoted to operators obeying $\hat{g}_n|n\rangle = g_n|n\rangle$ and $\hat{q}_n|n\rangle = q_n|n\rangle$, subscripts on operators denote the sector (n or m) on which they operate, and the arguments of $\hat{b}_n^-$ and $\hat{b}_m^-$ remind us that both are n -dependent.

Eqn. 64 makes clear that if not for the n -dependence of the operators the Hamiltonian $\hat{H}$ would be diagonalizable, or, equivalently, if g_n and q_n were constants the master equation would factorize into two individual birth-death processes. We may still, however, partition the full Hamiltonian as

$$\hat{H} = \hat{H}_0 + \hat{H}_1, \quad (69)$$

where $\hat{H}_0$ is a diagonalizable part (and $\hat{H}_1$ is the corresponding deviation from the diagonal form), and expand $|G\rangle$ in the eigenbasis of $\hat{H}_0$ to exploit the diagonality.

As with the multi-state system in Sec. IB, where we expand the solution in two different bases, there are several natural choices of eigenbasis of $\hat{H}_0$. Fig. 2 summarizes these choices diagrammatically: starting in the $|n, m\rangle$ basis (at the top of Fig. 2), one may expand in eigenfunctions either the first species, yielding the $|j, m\rangle$ basis (left), or the second species, yielding the $|n, k_n\rangle$ basis (right; in general we allow the parameter defining the second species' eigenfunctions to depend on n to reflect the regulation of the second species by the first. From either the $|j, m\rangle$ or the $|n, k_n\rangle$ basis, one may expand in eigenfunctions the remaining species, yielding either the $|j, k_n\rangle$ basis (bottom left), in which the second species' eigenfunctions depend on the first species' copy number

n , or the $|j, k_j\rangle$ basis (bottom right), in which the second species' eigenfunctions depend on the first species' eigenmode number j . Both bases reduce to the $|j, k\rangle$ basis (bottom center) when the parameter of the second species' eigenfunctions is a constant.

The $|n, m\rangle$ and $|j, m\rangle$ bases are less numerically useful than the other bases: as discussed above and detailed in Fig. 3, the $|n, m\rangle$ basis is numerically inefficient; and the $|j, m\rangle$ basis does not exploit the natural structure of the problem, since, unlike the other bases, it neither retains the tridiagonal structure in n nor gains a lower triangular structure in k (see the sections below). Each of the remaining bases, however, has preferable properties in terms of numerical stability and ability to represent the function sparsely yet accurately (either using a few values of n in the $|n, k_n\rangle$ basis or a few values of j in the $|j, k\rangle$, $|j, k_j\rangle$, or $|j, k_n\rangle$ bases, for example). Moreover, the equation of motion simplifies differently in each of the different bases. We present the derivations of the equations of motion in the following sections, beginning with the $|j, k\rangle$ basis, generalizing to the $|j, k_j\rangle$ basis, moving to the $|n, k_n\rangle$ basis, and ending with the $|j, k_n\rangle$ basis.

2. The $|j, k\rangle$ basis

For expository purposes we start by recalling the spectral representation used in previous work [34]. We choose the diagonal part of the Hamiltonian to correspond to two birth-death process with constant production rates $\bar{g}$ and $\bar{q}$,

$$\hat{H}_0 = \hat{b}_n^+ \bar{b}_n^- + \rho \hat{b}_m^+ \bar{b}_m^-, \quad (70)$$

where $\bar{b}_n^- = \hat{a}_n^- - \bar{g}$ and $\bar{b}_m^- = \hat{a}_m^- - \bar{q}$. As in Sec. IB, the parameters $\bar{g}$ and $\bar{q}$ act as gauge choices: their values can be set arbitrarily and thus can affect the numerical stability of the method. The nondiagonal part,

$$\hat{H}_1 = \hat{b}_n^+ \hat{\Gamma}_n + \rho \hat{b}_m^+ \hat{\Delta}_n, \quad (71)$$

captures the deviations $\hat{\Gamma}_n = \bar{g} - \hat{g}_n$ and $\hat{\Delta}_n = \bar{q} - \hat{q}_n$ of the regulation functions from the constant rates. We expand the generating function as

$$|G\rangle = \sum_{jk} G_{jk} |j, k\rangle, \quad (72)$$

where $|j, k\rangle$ is the eigenbasis of $\hat{H}_0$, i.e.

$$\hat{H}_0 |j, k\rangle = (j + \rho k) |j, k\rangle. \quad (73)$$

The eigenbasis is parameterized by the rates $\bar{g}$ and $\bar{q}$, meaning in position space $\langle x|j\rangle$ is as in Eqn. 38 and similarly for $\langle y|k\rangle$ with $x \rightarrow y$, $j \rightarrow k$, and $\bar{g} \rightarrow \bar{q}$.

With Eqns. 70-72, projecting the conjugate state $\langle j, k|$ onto Eqn. 63 yields

$$\begin{aligned} \dot{G}_{jk} = & -(j + \rho k) G_{jk} - \sum_{j'k'} \langle j | \hat{b}_n^+ \hat{\Gamma}_n | j' \rangle \langle k | k' \rangle G_{j'k'} \\ & - \rho \sum_{j'k'} \langle k | \hat{b}_m^+ | k' \rangle \langle j | \hat{\Delta}_n | j' \rangle G_{j'k'} \end{aligned} \quad (74)$$

(where the dummy indices j and k in Eqn. 72 have been changed to j' and k' respectively in Eqn. 74). Recalling Eqn. 24 and restricting attention to steady state, Eqn. 74 becomes

$$\begin{aligned} 0 = & -(j + \rho k) G_{jk} - \sum_{j'} \Gamma_{j-1, j'} G_{j'k} \\ & - \rho \sum_{j'} \Delta_{jj'} G_{j', k-1}, \end{aligned} \quad (75)$$

where the deviations have been rotated into the eigenbasis as

$$\Gamma_{jj'} = \langle j | \hat{\Gamma}_n | j' \rangle = \sum_n \langle j | n \rangle (\bar{g} - g_n) \langle n | j' \rangle, \quad (76)$$

$$\Delta_{jj'} = \langle j | \hat{\Delta}_n | j' \rangle = \sum_n \langle j | n \rangle (\bar{q} - q_n) \langle n | j' \rangle. \quad (77)$$

Eqn. 75 is subdiagonal in k and is therefore similar to Eqn. 43 in that the last term acts as a source term. It is initialized using

$$G_{j0} = \langle j, k=0 | G \rangle = \sum_{nm} p_{nm} \langle j | n \rangle \langle 0 | m \rangle \quad (78)$$

$$= \sum_n p_n \langle j | n \rangle, \quad (79)$$

(cf. Eqn. 26) with known p_n (cf. Eqn. 60), then solved at each subsequent k using the result for $k-1$. Eqn. 75 can be written in linear algebraic notation as

$$\vec{G}_k = -\rho (\mathbf{D}^k + \mathbf{S}^- \mathbf{\Gamma})^{-1} \mathbf{\Delta} \vec{G}_{k-1}, \quad (80)$$

where $\vec{G}_k$ is a vector over j , bold denotes matrices, and $D_{jj'}^k = (j + \rho k) \delta_{jj'}$ and $S_{jj'}^- = \delta_{j-1, j'}$ are diagonal and subdiagonal matrices, respectively. Eqn. 80 makes clear that the solution involves only matrix multiplication and the inversion of a J -by- J matrix K times, where J and K are cutoffs in the eigenmode numbers j and k , respectively. In fact, if the first species obeys a simple birth-death process, i.e. $g_n = g = \text{constant}$, setting $\bar{g} = g$ makes $\Gamma_{jj'} = 0$, and, since $\mathbf{D}^k$ is diagonal, the solution involves only matrix multiplication. The decomposition of the master equation into a linear algebraic equation results in huge gains in efficiency over direct solution in the $|n, m\rangle$ basis; the efficiency of all bases presented in this section is described in Sec. IIB and illustrated in Fig. 3.

Recalling Eqns. 62 and 72, the joint distribution is retrieved from G_{jk} via the inverse transform

$$p_{nm} = \sum_{jk} \langle n | j \rangle G_{jk} \langle m | k \rangle, \quad (81)$$

a computation again involving only matrix multiplication. The mixed product matrices $\langle n | j \rangle$ and $\langle m | k \rangle$ are computed as described in Appendix A.

3. The $|j, k_j\rangle$ basis

The $|j, k\rangle$ basis treats both genes similarly by expanding each around a constant production rate. We may instead imagine an eigenbasis that more closely reflects the underlying asymmetry imposed by the regulation, and make the basis of the second gene a function of that of the first. That is, we expand the first gene in a basis $|j\rangle$ with gauge $\bar{g}$ as before, but now we expand the second gene in a basis $|k_j\rangle$ with a j -dependent local gauge $\bar{q}_j$. We write the generating function as

$$|G\rangle = \sum_{jk} G_{jk} |j, k_j\rangle, \quad (82)$$

and Eqn. 63 at steady state becomes

$$0 = - \sum_{jk} G_{jk} \left[\hat{H}_0(j) + \hat{H}_1(j) \right] |j, k_j\rangle, \quad (83)$$

where we have partitioned the Hamiltonian for each j as

$$\hat{H}_0(j) = \hat{b}_n^+ \bar{b}_n^- + \rho \hat{b}_m^+ \bar{b}_m^-(j), \quad (84)$$

$$\hat{H}_1(j) = \hat{b}_n^+ \hat{\Gamma}_n + \rho \hat{b}_m^+ \hat{\Delta}_n(j), \quad (85)$$

with $\bar{b}_n^- = \hat{a}_n^- - \bar{g}$ and $\hat{\Gamma}_n = \bar{g} - \hat{g}_n$ as before and now $\bar{b}_m^-(j) = \hat{a}_m^- - \bar{q}_j$ and $\hat{\Delta}_n(j) = \bar{q}_j - \hat{q}_n$. Note that this basis enjoys the eigenvalue equation

$$\hat{H}_0(j) |j, k_j\rangle = (j + \rho k) |j, k_j\rangle. \quad (86)$$

Projecting the conjugate state $\langle j, k_j|$ onto Eqn. 83 yields, after some simplification (cf. Appendix C), the equation of motion

$$0 = (j + \rho k) G_{jk} + \sum_{j'} \Gamma_{j-1, j'} G_{j'k} + \sum_{\ell=1}^k \sum_{j'} \left(\Gamma_{j-1, j'} V_{jj'}^\ell + \rho \Delta_{jj'} V_{jj'}^{\ell-1} \right) G_{j', k-\ell}, \quad (87)$$

where $\Gamma_{jj'}$ is as in Eqn. 76 and

$$\Delta_{jj'} = \langle j | \hat{\Delta}_n(j) | j' \rangle = \sum_n \langle j | n \rangle (\bar{q}_j - q_n) \langle n | j' \rangle, \quad (88)$$

$$V_{jj'}^\ell = \frac{(-Q_{jj'})^\ell}{\ell!}, \quad (89)$$

with $Q_{jj'} = \bar{q}_j - \bar{q}_{j'}$. Eqn. 87 can be written linear algebraically as

$$\vec{G}_k = -(\mathbf{D}^k + \mathbf{S}^- \mathbf{\Gamma})^{-1} \times \sum_{\ell=1}^k [(\mathbf{S}^- \mathbf{\Gamma}) * \mathbf{V}^\ell + \rho \mathbf{\Delta} * \mathbf{V}^{\ell-1}] \vec{G}_{k-\ell}, \quad (90)$$

where $D_{jj'}^k$ and $S_{jj'}^-$ are defined as before (cf. Eqn. 80), and $*$ denotes an element-by-element matrix product. Once again, Eqn. 90 is lower-triangular in k and requires

only matrix multiplication and the inversion of a J -by- J matrix K times. $\vec{G}_k$ is initialized as in Eqn. 79, and the k th term is computed from the previous $k' < k$ terms. The joint distribution is retrieved via the inverse transform

$$p_{nm} = \sum_{jk} \langle n | j \rangle G^{jk} \langle m | k_j \rangle. \quad (91)$$

4. The $|n, k_n\rangle$ basis

Expansion in either the $|j, k\rangle$ or the $|j, k_j\rangle$ basis conveniently turns the master equation into a lower-triangular linear algebraic equation and replaces cutoffs in protein number with cutoffs in eigenmode number (which can be smaller with appropriate choices of gauge). However, these bases sacrifice the original tridiagonal structure of the master equation in the copy number of first gene, n . Therefore we now consider a mixed representation, in which the first gene remains in protein number space $|n\rangle$, and we expand the second gene in an n -dependent eigenbasis $|k_n\rangle$,

$$|G\rangle = \sum_{nk} G_{nk} |n, k_n\rangle. \quad (92)$$

If the rate parameter of the $|k_n\rangle$ basis were the regulation function q_n , it would be natural to make $\hat{H}_0$ the entire Hamiltonian, Eqn. 64. For generality we will instead allow the rate parameter of the $|k_n\rangle$ basis to be an arbitrary n -dependent local gauge $\bar{q}_n$, such that Eqn. 63 at steady state naturally partitions as

$$0 = - \sum_{nk} G_{nk} \left[\hat{H}_0(n) + \hat{H}_1(n) \right] |n, k_n\rangle, \quad (93)$$

where

$$\hat{H}_0(n) = \hat{b}_n^+ \bar{b}_n^- + \rho \hat{b}_m^+ \bar{b}_m^-(n), \quad (94)$$

$$\hat{H}_1(n) = \rho \hat{b}_m^+ \hat{\Delta}_n(n), \quad (95)$$

with $\bar{b}_m^-(n) = \hat{a}_m^- - \bar{q}_n$ and $\hat{\Delta}_n(n) = \bar{q}_n - \hat{q}_n$. Note that $|n, k_n\rangle$ is not the eigenbasis of $\hat{H}_0(n)$, but rather $\hat{H}_0(n) |n, k_n\rangle$ retains the original tridiagonal structure in n , i.e.

$$\begin{aligned} \hat{H}_0(n) |n, k_n\rangle &= (g_n + n) |n, k_n\rangle - g_n |n+1, k_n\rangle \\ &\quad - n |n-1, k_n\rangle + \rho k |n, k_n\rangle, \end{aligned} \quad (96)$$

where Eqns. 65, 67, 12, and 13 are recalled in applying the first term of $\hat{H}_0(n)$.

Projecting the conjugate state $\langle n, k_n|$ onto Eqn. 93 yields, after some simplification (cf. Appendix C), the equation of motion

$$\begin{aligned} &g_{n-1} G_{n-1, k} + (n+1) G_{n+1, k} - (g_n + n + \rho k) G_{nk} \\ &= -g_{n-1} \sum_{\ell=1}^k V_{n\ell}^- G_{n-1, k-\ell} - (n+1) \sum_{\ell=1}^k V_{n\ell}^+ G_{n+1, k-\ell} \\ &\quad + \rho \Delta_n G_{n, k-1}, \end{aligned} \quad (97)$$

where

$$\Delta_n = \bar{q}_n - q_n, \quad (98)$$

$$V_{n\ell}^\pm = \frac{(-Q_n^\pm)^\ell}{\ell!}, \quad (99)$$

and $Q_n^\pm = \bar{q}_n - \bar{q}_{n\pm 1}$. Eqn. 97 can be written linear algebraically as

$$\begin{aligned} \vec{G}_k = & (\mathbf{T}^k)^{-1} \left\{ (\mathbf{S}^- \vec{g}) * \text{diag}[\mathbf{V}^- (\mathbf{S}^- \tilde{\mathbf{G}})^T] \right. \\ & + (\mathbf{S}^+ \vec{n}) * \text{diag}[\mathbf{V}^+ (\mathbf{S}^+ \tilde{\mathbf{G}})^T] \\ & \left. + \rho \vec{\Delta} * \vec{G}_{k-1} \right\}, \end{aligned} \quad (100)$$

where $\mathbf{X}^T$ indicates the transpose of $\mathbf{X}$, $*$ denotes an element-by-element product, $S_{nn'}^\pm = \delta_{n\pm 1, n'}$ are super- (+) and subdiagonal (-) matrices, $T_{nn'}^k = g_{n-1} \delta_{n-1, n'} + (n+1) \delta_{n+1, n'} - (g_n + n + \rho k) \delta_{nn'}$ is a tridiagonal matrix, and $\tilde{\mathbf{G}}$, which is $\mathbf{G}$ with the columns reversed (i.e. $\tilde{G}_{n\ell} = G_{n, k-\ell}$), is built incrementally in k . As with the $|j, k\rangle$ and $|j, k_j\rangle$ bases, the k th term is slaved to the previous $k' < k$ terms; in total the solution requires K inversions of an N -by- N matrix. However, here the task of inversion is simplified because the matrix to be inverted is tridiagonal. In fact, using the Thomas algorithm [40], we obtain an analytic solution for the case of constant production of the first gene and threshold regulation of the second gene, as described in section II C.

The solution is initialized at $k = 0$ using

$$\begin{aligned} G_{n0} &= \langle n, (k=0)_n | G \rangle = \sum_{n'm} \langle n | n' \rangle \langle 0_n | m \rangle p_{n'm} \\ &= p_n \end{aligned} \quad (101)$$

(cf. Eqn. 26), where p_n is known (cf. Eqn. 60), and the joint distribution is retrieved via the inverse transform

$$p_{nm} = \langle n, m | G \rangle = \sum_k G_{nk} \langle m | k_n \rangle. \quad (102)$$

5. The $|j, k_n\rangle$ basis

We now consider a basis which employs both the constant-rate eigenfunctions $|j\rangle$ in the n sector and the n -dependent eigenfunctions $|k_n\rangle$ in the m sector. Expressing the joint distribution directly in terms of the eigenbasis expansion of the generating function [51], we write

$$p_{nm} = \sum_{jk} G_{jk} \langle n, m | j, k_n \rangle, \quad (103)$$

where, as in the $|j, k\rangle$ and $|j, k_j\rangle$ bases, the $|j\rangle$ are parameterized by the constant rate $\bar{g}$, and, as in the $|n, k_n\rangle$ basis, the $|k_n\rangle$ are parameterized by the arbitrary function $\bar{q}_n$. The inverse of Eqn. 103 is

$$G_{jk} = \sum_{nm} p_{nm} \langle j, k_n | n, m \rangle. \quad (104)$$

Substituting Eqn. 103 into Eqn. 58 at steady state gives, after some simplification (cf. Appendix C),

$$\begin{aligned} 0 = & \sum_{jk} G_{jk} \langle n, m | \{-\hat{H} | j, k_n \rangle \\ & + \hat{a}_n^+ \hat{g}_n | j, \delta_- k_n \rangle + \hat{a}_n^- | j, \delta_+ k_n \rangle \}, \end{aligned} \quad (105)$$

where $\hat{H}$ is defined as in Eqn. 64, $\hat{a}_n^+$ and $\hat{a}_n^-$ act as in Eqns. 12-15, and

$$|\delta_\pm k_n\rangle = |k_{n\pm 1}\rangle - |k_n\rangle. \quad (106)$$

We may now partition $\hat{H} = \hat{H}_0 + \hat{H}_1$ as

$$\hat{H}_0(n) = \hat{b}_n^+ \bar{b}_n^- + \rho \hat{b}_m^+ \bar{b}_m^-(n), \quad (107)$$

$$\hat{H}_1(n) = \hat{b}_n^+ \hat{\Gamma}_n + \rho \hat{b}_m^+ \hat{\Delta}_n(n), \quad (108)$$

with $\bar{b}_n^- = \hat{a}_n^- - \bar{g}$ and $\hat{\Gamma}_n = \bar{g} - \hat{g}_n$ as in the $|j, k\rangle$ and $|j, k_j\rangle$ bases, and $\bar{b}_m^-(n) = \hat{a}_m^- - \bar{q}_n$ and $\hat{\Delta}_n(n) = \bar{q}_n - \hat{q}_n$ as in the $|n, k_n\rangle$ basis. Noting that $|j, k_n\rangle$ is the eigenbasis of $\hat{H}_0(n)$, i.e.

$$\hat{H}_0(n) |j, k_n\rangle = (j + \rho k) |j, k_n\rangle, \quad (109)$$

Eqn. 105 becomes, after more simplification (cf. Appendix C),

$$\begin{aligned} 0 = & -(j + \rho k) G_{j'k} - \sum_{j'} \Gamma_{j-1, j'} G_{j'k} - \rho \sum_{j'} \Delta_{jj'} G_{j', k-1} \\ & + \sum_{\pm} \sum_{j'} \sum_{\ell=1}^k \Lambda_{jj'}^{\pm \ell} G_{j', k-\ell}, \end{aligned} \quad (110)$$

where $\Gamma_{jj'}$ is as in Eqn. 76,

$$\Delta_{jj'} = \sum_n \langle j | n \rangle (\bar{q}_n - q_n) \langle n | j' \rangle, \quad (111)$$

$$\Lambda_{jj'}^{+ \ell} = \sum_n \langle j | n \rangle (n+1) \langle n+1 | j' \rangle V_{nl}^+, \quad (112)$$

$$\Lambda_{jj'}^{- \ell} = \sum_n \langle j | n \rangle g_{n-1} \langle n-1 | j' \rangle V_{nl}^-, \quad (113)$$

and $V_{n\ell}^\pm$ is as in Eqn. 99. Linear algebraically,

$$\begin{aligned} \vec{G}_k = & (\mathbf{D}^k + \mathbf{S}^- \mathbf{\Gamma})^{-1} \\ & \times \left(-\rho \mathbf{\Delta} \vec{G}_{k-1} + \sum_{\pm} \sum_{\ell=1}^k \mathbf{\Lambda}^{\pm \ell} \vec{G}_{k-\ell} \right), \end{aligned} \quad (114)$$

with $D_{jj'}^k$ and $S_{jj'}^-$ defined as before (cf. Eqn. 80), revealing once again a lower-triangular equation (i.e. each k th term is slaved to the previous $k' < k$ terms) requiring only matrix multiplication and the inversion of a J -by- J matrix K times. Recalling Eqn. 104, the scheme is initialized using

$$G_{j0} = \sum_{nm} p_{nm} \langle j | n \rangle \langle (k=0)_n | m \rangle = \sum_n p_n \langle j | n \rangle \quad (115)$$

(cf. Eqn. 26) with known p_n (cf. Eqn. 60), and the joint distribution is retrieved using Eqn. 103.

B. Comparison of the representations

The spectral representations in Sec. II A produce equations of motion with similar levels of numerical complexity. In all cases, the original two-dimensional master equation has been reduced by the lower triangular structure in the second gene's eigenmode number k to a hierarchy of evaluations of one dimensional problems. The bases differ in the rate parameters, or equivalently gauge freedoms, that one is free to choose: the $|j, k\rangle$ basis requires two constants $\bar{g}$ and $\bar{q}$; the $|j, k_j\rangle$ basis requires $\bar{g}$ and a J -valued vector $\bar{q}_j$; the $|n, k_n\rangle$ basis requires a N -valued vector $\bar{q}_n$; and the $|j, k_n\rangle$ basis requires $\bar{g}$ and $\bar{q}_n$.

The bases also differ in the types of problems for which they are most suitable. For example, the $|j, k\rangle$, $|j, k_j\rangle$, and $|j, k_n\rangle$ bases, which all expand the parent species in eigenfunctions $|j\rangle$, are best when a cutoff in j is most appropriate, such as when the parent distribution is a Poisson. The $|n, k_n\rangle$ basis, on the other hand, is useful when a cutoff is n is most appropriate, such as when the parent species is concentrated at low protein number. Different bases are more robust to numerical errors for different regulation functions as well: the $|n, k_n\rangle$ and $|j, k_n\rangle$ bases, which both rely upon repeated manipulation of the object $Q_n^\pm = \bar{q}_n - \bar{q}_{n\pm 1}$, are best for smooth regulation functions, for which the differences between $\bar{q}_n$ and $\bar{q}_{n+1}$ are small; the $|j, k\rangle$ and $|j, k_j\rangle$ bases on the other hand, which involve the deviations $\bar{q} - q_n$ and $\bar{q}_j - q_n$ respectively, are less susceptible to numerical error given sharp regulation functions, such as a threshold.

As indicated in Figure 2, the $|j, k\rangle$ basis can be viewed as a special case of either the $|j, k_j\rangle$ basis with $\bar{q}_j = \bar{q}$ (in which case Eqn. 87 reduces to Eqn. 75) or of the $|j, k_n\rangle$ basis with $\bar{q}_n = \bar{q}$ (in which case Eqn. 110 reduces to Eqn. 75). Although possible in principle, expanding in the $|j, m\rangle$ basis does not exploit the natural structure of the problem, since it neither retains the tridiagonal structure in n nor gains the lower triangular structure in k . This example explicitly shows that not all bases are good candidates for simplifying the master equation.

The strength of all the spectral bases discussed in this section, and of the proposed spectral method in general, is that it allows for a fast and accurate calculation of full steady state probability distributions of the number of protein molecules in a gene regulatory network. In Fig. 3 we demonstrate this property for the two-gene system by plotting error versus computational runtime for each spectral basis, as well as for a stochastic simulation using a varying step Monte Carlo procedure [32]. For error we use the Jensen-Shannon divergence [41] (a measure in bits between two probability distributions) between the distribution p_{nm} computed in the $|n, m\rangle$ basis (via iterative solution of the original master equation) and the distribution computed either via the spectral formulae in this section or by stochastic simulation. We plot this measure against the runtime of each method, scaled by the runtime of the iterative solution in the $|n, m\rangle$ basis (all




FIG. 3: Error vs. runtime for the spectral method and stochastic simulation. Error is the Jensen-Shannon divergence [41] between p_{nm} obtained using the $|n, m\rangle$ basis (via iterative solution of the original master equation) and that obtained using the $|j, k\rangle$ basis (circles; cf. Eqn. 80), the $|j, k_j\rangle$ basis (triangles; cf. Eqn. 90), the $|n, k_n\rangle$ basis (squares; cf. Eqn. 100), the $|j, k_n\rangle$ basis (diamonds; cf. Eqn. 114), or stochastic simulation [32] (dots). Runtimes are scaled by that of the iterative solution, 150 seconds (in MATLAB). Spectral basis data is obtained by varying K , the cutoff in the eigenmode number k of the second gene; simulation data is obtained by varying the integration time. The input distribution $p_n = \pi_1 e^{-\lambda_1} \lambda_1^n / n! + (1 - \pi_1) e^{-\lambda_2} \lambda_2^n / n!$ (from which g_n is calculated via Eqn. 59) is a mixture of two Poisson distributions with $\lambda_1 = 0.5$, $\lambda_2 = 15$, and $\pi_1 = 0.5$. The regulation function $q_n = q_- + (q_+ - q_-) n^\nu / (n^\nu + n_0^\nu)$ is a Hill function with $q_- = 1$, $q_+ = 11$, $n_0 = 7$, and $\nu = 2$. The gauge choices used (cf. Fig. 2) are $\bar{g} = \sum_n p_n g_n$, $\bar{q} = \sum_n p_n q_n$, $\bar{q}_n = q_n$, and $\bar{q}_j = \sum_n \langle j | n \rangle q_n \langle n | j \rangle$. The cutoffs used are $J = 80$ for the eigenmode number j of the first gene and $N = 50$ for the protein numbers n and m . **Inset:** The joint probability distribution p_{nm} . The peak at low protein number extends to $p_{00} \approx 0.1$.

numerical experiments are performed in MATLAB). We find that the computations via the spectral bases achieve accuracy up to machine precision $\sim 10^3$ - 10^4 times faster than the iterative method's runtime and $\sim 10^7$ - 10^8 times faster than the runtime necessary for the stochastic simulation to achieve the same accuracy. Computation in the $|j, k\rangle$ basis is most efficient since its equation of motion is simplest (cf. Eqn. 80); the $|j, k_j\rangle$ and $|j, k_n\rangle$ bases tend to be slightly less efficient since they require inner loops over ℓ (cf. Eqns. 90 and 114). Fig. 3 demonstrates the tremendous gain in performance achieved by the joint analytic-numerical spectral method over traditional simulation approaches.

C. An analytic solution

In general, the equations of motion in the spectral representations (Eqns. 75, 87, 97, and 110) need to be evaluated numerically. In the case of the $|n, k_n\rangle$ basis, however, we can exploit the tridiagonal structure of Eqn. 97 to find an exact analytic solution. Specifically, in the case of a Poisson parent ($g_n = g = \text{constant}$) and for threshold regulation, i.e.

$$q_n = \begin{cases} q_- & \text{for } n \leq n_0 \\ q_+ & \text{for } n > n_0, \end{cases} \quad (116)$$

setting $\bar{q}_n = q_n$ makes Eqn. 97

$$gG_{n-1,k} + (n+1)G_{n+1,k} - (\rho k + g + n)G_{nk} = -g\phi_k^- \delta_{nn_1} - n_1\phi_k^+ \delta_{nn_0}, \quad (117)$$

where

$$\phi_k^- = \sum_{\ell=1}^k \frac{(-\Delta)^\ell}{\ell!} G_{n_0, k-\ell}, \quad (118)$$

$$\phi_k^+ = \sum_{\ell=1}^k \frac{\Delta^\ell}{\ell!} G_{n_1, k-\ell}, \quad (119)$$

$\Delta = q_+ - q_-$, and $n_1 = n_0 + 1$. Eqn. 117 is solved using the tridiagonal matrix algorithm (also called the Thomas algorithm [40]), as described in detail in Appendix D. The result is an analytic expression for the k th column of G_{nk} in terms of its previous columns (i.e. the matrix inversion has been performed explicitly),

$$G_{nk} = \frac{n_1}{g} \frac{(n_0 - 1)!}{n!} \frac{\eta_n^k}{\eta_{n_0-1}^k} \times \begin{cases} (\phi_k^+ + f_k F_{n_1}^k) / (\epsilon_{n_0}^k - 1) & n \leq n_0 \\ f_k F_n^k / \prod_{i=n_0}^{n-1} (\epsilon_i^k - 1) & n > n_0, \end{cases} \quad (120)$$

where

$$f_k = \phi_k^+ - \frac{gn_0}{n_1} \frac{\eta_{n_0-1}^k}{\eta_{n_0}^k} (\epsilon_{n_0}^k - 1) \phi_k^-, \quad (121)$$

$$F_n^k = \sum_{i=0}^{N-n} \prod_{\ell=n}^{n+i} \frac{1}{\epsilon_\ell^k - 1}, \quad (122)$$

$$\epsilon_n^k = \frac{\rho k + g + n}{gn} \frac{\eta_n^k}{\eta_{n-1}^k}, \quad (123)$$

$$\eta_n^k = \sum_{i=0}^n \frac{n!}{i!(n-i)!} g^{n-i} \prod_{\ell=0}^{i-1} (\rho k + \ell) \quad (124)$$

$$= \frac{1}{\Gamma(\rho k)} \int_0^t dt e^{-t} t^{(\rho k - 1)} (g + t)^n, \quad (125)$$

and N is the cutoff in protein number n . Along with the analytic form of the mixed product

$$\langle m | k_n \rangle = (-1)^k e^{-q_n} q_n^m k! \times \sum_{\ell=0}^{\min(m, k)} \frac{1}{\ell! (m - \ell)! (k - \ell)! (-q_n)^\ell} \quad (126)$$

(cf. Appendix A), Eqn. 120 in the limit $N \rightarrow \infty$ constitutes an exact analytic solution for the joint distribution p_{nm} , as calculated using Eqn. 102.

D. The threshold-regulated gene approximates the on/off gene

If a gene is regulated via a threshold function (cf. Eqn. 116), its steady state protein distribution p_m can be well approximated by the two-state process discussed in Sec. IC. To make the connection clear, we first observe that the off-state ($z = -$) corresponds to the first gene expressing the same or fewer proteins n than the threshold n_0 , i.e.

$$p_m^- = \sum_{n \leq n_0} p_{nm}, \quad (127)$$

and the on-state ($z = +$) corresponds to the first gene expressing more proteins than the threshold, i.e.

$$p_m^+ = \sum_{n > n_0} p_{nm}. \quad (128)$$

The dynamics of $p_m^\pm$ are then obtained by summing the master equation for two-gene regulation, Eqn. 58, over either all $n \leq n_0$ or all $n > n_0$, giving

$$\dot{p}_m^\pm = \rho [q_\pm p_{m-1}^\pm + (m+1)p_{m+1}^\pm - (q_\pm + m)p_m^\pm] \mp n_1 p_{n_1 m} \pm g_{n_0} p_{n_0 m}, \quad (129)$$

where $n_1 = n_0 + 1$. Making the approximations

$$\frac{p_m^-}{\pi_-} = p(m|-) \approx p(m|n_0) = \frac{p_{n_0 m}}{p_{n_0}}, \quad (130)$$

$$\frac{p_m^+}{\pi_+} = p(m|+) \approx p(m|n_1) = \frac{p_{n_1 m}}{p_{n_1}}, \quad (131)$$

where

$$\pi_- = \sum_m p_m^- = \sum_{n \leq n_0} p_n, \quad (132)$$

$$\pi_+ = \sum_m p_m^+ = \sum_{n > n_0} p_n \quad (133)$$

are the total probabilities of being in the off- and on-states respectively, and noting from Eqn. 59 that $g_{n_0} = n_1 p_{n_1} / p_{n_0}$, Eqn. 129 at steady state becomes

$$0 = q_z p_{m-1}^z + (m+1)p_{m+1}^z - (q_z + m)p_m^z + \sum_{z'} \Omega_{zz'} p_m^{z'}, \quad (134)$$

with $z = \pm$ and

$$\Omega = \begin{pmatrix} -\omega_+ & \omega_- \\ \omega_+ & -\omega_- \end{pmatrix}, \quad (135)$$

where

$$\omega_{\pm} = \frac{n_1 p_{n_1}}{\rho \pi_{\mp}}. \quad (136)$$

Eqns. 134-135 have the same form as Eqns. 1 and 47 at steady state with $n \rightarrow m$ and $g \rightarrow q$, and Eqn. 136 relates the effective switching rates $\omega_{\pm}$ to input and regulation parameters p_{n_1} , $\pi_{\pm}$, and n_1 , and the ratio ρ of the degradation rate of the second gene to that of the first. Note that Eqn. 136 satisfies

$$\frac{\pi_{-}}{\pi_{+}} = \frac{\omega_{-}}{\omega_{+}}, \quad (137)$$

in agreement with Eqn. 48, and exhibits the intuitive behavior that increasing ρ (i.e. decreasing the relative response rate of the first gene) is equivalent to decreasing the switching rates $\omega_{\pm}$.

A comparison of the distributions of a threshold-regulated gene with those of an on/off gene for various parameter settings reveals that Eqns. 130-131 are a good approximation. Fig. 4 shows a demonstration for a threshold-regulated system with a Poisson input distribution. In the first column, the mean g of the input lies above the threshold n_0 , making the output more likely to be in the on-state, i.e. $\pi_{+} > \pi_{-}$; in the second column, $g < n_0$, making $\pi_{+} < \pi_{-}$. In the first row $\rho < 1$; in the second row $\rho > 1$, corresponding to lower effective switching rates $\omega_{\pm}$ and producing bimodal distributions with peaks near the on/off rates $q_{\pm}$. In all examples, the approximation as a two-state process with switching rates given by Eqn. 136 agrees well with the actual output from threshold regulation.

III. REGULATION WITH BURSTS

The final system we consider combines the multi-state process used to model bursts of expression in Sec. I with gene regulation as discussed in Sec. II. Specifically we consider a system of two species, with protein numbers n and m , existing in Z possible states, distinguished by the settings of the two production rates g_z and q_z respectively, where $1 \leq z \leq Z$. Regulation is achieved by allowing the rates of transition among states affecting the production of the second gene to depend on the number n of proteins expressed by the first gene. Recalling Eqns. 1 and 58, the master equation describing the evolution of the joint probability distribution p_{nm}^z reads

$$\begin{aligned} \dot{p}_{nm}^z = & g_z p_{n-1,m}^z + (n+1) p_{n+1,m}^z - (g_z + n) p_{nm}^z \\ & + \rho [q_z p_{n,m-1}^z + (m+1) p_{n,m+1}^z - (q_z + m) p_{nm}^z] \\ & + \sum_{z'} \Omega_{zz'}(n) p_{nm}^{z'}, \end{aligned} \quad (138)$$

where the dependence of the stochastic matrix $\Omega_{zz'}$ on n incorporates the regulation.

As with the previously discussed models, Eqn. 138 benefits from spectral expansion, and for simplicity we




FIG. 4: Protein distributions for a gene regulated by a threshold function (dots; calculated via Eqn. 75) and a gene with two stochastic states (circles; calculated via Eqn. 43). The relationship between regulation parameters and state transition rates is given by Eqn. 136. In all panels the input to the regulation is a Poisson distribution with mean $g = 7$, and the regulation rates (cf. Eqn. 116) are $q_- = 2$ and $q_+ = 15$. In the first column the threshold is $n_0 = 4$ making $\pi_+ = 0.827 > \pi_- = 0.173$; in the second column $n_0 = 8$ making $\pi_+ = 0.271 < \pi_- = 0.729$. In the first row the ratio of the second gene's degradation rate to that of the first is $\rho = 0.1$; in the second row $\rho = 10$.

present only the formulation in the $|j, k\rangle$ basis, parameterized by constant rates $\bar{g}$ and $\bar{q}$ respectively, as in Secs. IB and IIA 2. As before the first step is to define the generating function

$$|G_z\rangle = \sum_{nm} p_{nm}^z |n, m\rangle, \quad (139)$$

with which Eqn. 138, upon summing over n and m against $|n, m\rangle$, becomes

$$|\dot{G}_z\rangle = -\hat{H}_z |G_z\rangle + \sum_{z'} \hat{\Omega}_{zz'} |G_{z'}\rangle, \quad (140)$$

where

$$\hat{H}_z = \hat{b}_n^+ \hat{b}_{nz}^- + \rho \hat{b}_m^+ \hat{b}_{mz}^- \quad (141)$$

$$\hat{b}_n^+ = \hat{a}_n^+ - 1, \quad (142)$$

$$\hat{b}_m^+ = \hat{a}_m^+ - 1, \quad (143)$$

$$\hat{b}_{nz}^- = \hat{a}_n^- - g_z, \quad (144)$$

$$\hat{b}_{mz}^- = \hat{a}_m^- - q_z, \quad (145)$$

and $\hat{\Omega}_{zz'}$ is $\Omega_{zz'}(n)$ with every instance of n replaced by the number operator $\hat{a}_n^+ \hat{a}_n^-$. Defining $\bar{b}_n^- = \hat{a}_n^- - \bar{g}$ and $\bar{b}_m^- = \hat{a}_m^- - \bar{q}$, we partition the Hamiltonian as $\hat{H}_z = \hat{H}_0 + \hat{H}_1^z$, with

$$\hat{H}_0 = \hat{b}_n^+ \bar{b}_n^- + \rho \hat{b}_m^+ \bar{b}_m^- \quad (146)$$

the operator of which $|j, k\rangle$ is the eigenbasis, i.e.

$$\hat{H}_0|j, k\rangle = (j + \rho k)|j, k\rangle, \quad (147)$$

and

$$\hat{H}_1^z = \hat{b}_n^+ \Gamma_z + \rho \hat{b}_m^+ \Delta_z \quad (148)$$

capturing the deviations $\Gamma_z = \bar{g} - g_z$ and $\Delta_z = \bar{q} - q_z$ of the constant rates from the state-dependent rates. Upon expanding the generating function in the eigenbasis,

$$|G_z\rangle = \sum_{jk} G_{jk}^z |j, k\rangle, \quad (149)$$

and taking dummy indices $j \rightarrow j'$ and $k \rightarrow k'$, projecting the conjugate state $\langle j, k|$ onto Eqn. 140 gives

$$\begin{aligned} \dot{G}_{jk}^z &= -(j + \rho k)G_{jk}^z - \Gamma_z G_{j-1,k}^z - \Delta_z G_{j,k-1}^z \\ &\quad + \sum_{z'} \sum_{j'} \langle j | \hat{\Omega}_{zz'} | j' \rangle G_{j'k}^{z'}, \end{aligned} \quad (150)$$

where the components of $\hat{\Omega}_{zz'}$ need only be evaluated in the j sector, not the k sector, because the transition rates depend on only n , not m (cf. Eqn. 138). Like Eqns. 43 and 75, Eqn. 150 is subdiagonal in k and thus far more efficient to solve than the original master equation, Eqn. 138, as we demonstrate for a special case in the next section. The joint distribution is retrieved from G_{jk}^z via inverse transform,

$$p_{nm}^z = \sum_{jk} \langle n | j \rangle G_{jk}^z \langle m | k \rangle, \quad (151)$$

with the mixed products calculated as in Appendix A.

A. The four-state process

As a simple example of the model in Eqn. 138, we consider a system in which each of the two species has an on-state and an off-state, and the transition rate of the second species to its on-state is a function of the number of copies of the first species. This system models both (i) a single gene for which the production of proteins depends on the number of transcripts, and each is produced in on- and off-states by the binding and unbinding of ribosomes and RNA polymerase respectively, and (ii) one gene regulating another with each undergoing burst-like expression.

There are a total of $Z = 4$ states, i.e.

$$p_{nm}^z = (p_{nm}^{--}, p_{nm}^{+-}, p_{nm}^{-+}, p_{nm}^{++}). \quad (152)$$

where the first signed index denotes the state of the first gene (with protein count n) and the second signed index denotes the state of the second gene (with protein count m). Defining $g_{\pm}$ as the production rates of the

first species in its on- (+) and off-states (-), and similarly $q_{\pm}$ for the second species, the production rates of the $Z = 4$ states are:

$$g_z = (g_-, g_+, g_-, g_+), \quad (153)$$

$$q_z = (q_-, q_-, q_+, q_+). \quad (154)$$

Defining $\omega_{\pm}$ as the transition rates of the first species to (+) and from (-) its on-state, and similarly $\alpha_{\pm}$ for the second species, the transition matrix takes the form

$$\Omega_{zz'}(n) = \begin{pmatrix} -\omega_+ - \alpha_+(n) & \omega_- & \alpha_- & 0 \\ \omega_+ & -\omega_- - \alpha_+(n) & 0 & \alpha_- \\ \alpha_+(n) & 0 & -\omega_+ - \alpha_- & \omega_- \\ 0 & \alpha_+(n) & \omega_+ & -\omega_- - \alpha_- \end{pmatrix}. \quad (155)$$

The simple form $\alpha_+(n) = cn^{\nu}$ for constant c and integer ν corresponds to the first species activating the second as a multimer, with ν the order of the multimerization. In the limit of fast switching this description reduces to a Hill function with cooperativity ν [28]. Recalling that $\hat{b}_n^+ = \hat{a}_n^+ - 1$ and $\bar{b}_n^- = \hat{a}_n^- - \bar{g}$, the n -dependent terms of $\langle j | \hat{\Omega}_{zz'} | j' \rangle$ are evaluated as

$$\langle j | \alpha_+ (\hat{a}_n^+ \hat{a}_n^-) | j' \rangle = c \langle j | [(\hat{b}_n^+ + 1)(\bar{b}_n^- + \bar{g})]^{\nu} | j' \rangle, \quad (156)$$

Since $\hat{b}_n^+$ and $\bar{b}_n^-$ raise and lower $|j'\rangle$ states respectively (cf. Eqns. 22-23), the modified transition matrix $\langle j | \hat{\Omega}_{zz'} | j' \rangle$ is nearly diagonal, with nonzero terms only for $|j - j'| \leq \nu$.

Eqn. 150 at steady state,

$$\begin{aligned} &-(j + \rho k)G_{jk}^z - \Gamma_z G_{j-1,k}^z + \sum_{z'} \sum_{j'} \langle j | \hat{\Omega}_{zz'} | j' \rangle G_{j'k}^{z'} \\ &= \Delta_z G_{j,k-1}^z, \end{aligned} \quad (157)$$

is solved successively in k , requiring the inversion of a $4J$ -by- $4J$ matrix K times. It is initialized at $k = 0$ by computing the null space of the left hand side and normalizing with $\sum_z G_{00}^z = 1$ (cf. Eqn. 35). The joint distribution p_{nm}^z is retrieved via inverse transform (Eqn. 151).

With $\nu = 2$, a typical solution of Eqn. 157 takes a few seconds (in MATLAB), which, depending on the cutoff N , is $\sim 10^2$ - 10^3 times faster than direct solution of the master equation, Eqn. 138, by iteration, for equivalent accuracy. The advantage of such a large efficiency gain is that it allows repeated evaluations of the governing equation, necessary for parameter inference or optimization [34]. We demonstrate this possibility in the next section by finding and interpreting the solutions that optimize the information flow from the first to the second species.

B. The information-optimal solution

Cells use regulatory processes to transmit relevant information from one species to the next [42, 43, 44, 45, 46]. Information processing is quantified by the mutual information I , which, between the first and second species in the four-state process, is

$$I = \sum_{nm} p_{nm} \log_2 \frac{p_{nm}}{p_n p_m}, \quad (158)$$

where the distributions p_{nm} , p_n , and p_m are obtained from summing the joint distribution p_{nm}^z (cf. Eqn. 151), and the log is taken with base 2 to give I in bits.

Upon optimization of I for the four-state process, two distinct types of optimal solutions become clear: those in which the distribution p_{nm} has one peak, and those in which p_{nm} has two peaks. The former occur when copy number is constrained to be low, and switching rates are constrained to be near the decay rates of both species, producing a single peak at low copy number (see lower left inset of Fig. 5B). As these constraints are lifted, it is optimal for the switching rates of the parent species to become much less than the decay rate. The slow switching produces a second peak whose location is specified by the on-rate of each species (see upper right inset of Fig. 5B).

To quantify the transition between the two types of solutions, we numerically optimized mutual information over parameters g_+ , q_+ , ω_- , ω_+ , α_- , and c (the off-rates g_- and q_- were fixed at 0; the cooperativity ν was fixed at 2; and the decay rate ratio ρ was fixed at 1). Information may always be trivially optimized by allowing infinite copy number or arbitrary separation of relevant timescales. We limit copy number by constraining the gain

$$\gamma = \frac{\Gamma + \Delta}{2}, \quad (159)$$

defined as the average of the parent gain $\Gamma = g_+ - g_-$ and the child gain $\Delta = q_+ - q_-$. Since $g_- = q_- = 0$, the maximum number of particles is dictated by the on-rates g_+ and q_+ , and thus constraining γ limits the copy number. We limit separation between the switching timescales and the decay timescales by constraining the stiffness

$$\sigma = \frac{1}{4} (|\log_{10} \omega_-| + |\log_{10} \omega_+| + |\log_{10} \alpha_-| + |\log_{10} [\alpha_+ \langle n^\nu \rangle]|), \quad (160)$$

where the average $\langle n^\nu \rangle$ is taken over p_n . Stiffness σ is the average of the absolute deviation of (the logs of) all four switching rates from the (unit) decay rates, so constraining σ prevents fast or slow switching. Gain is fixed by varying g_+ and q_+ such that γ is a constant, and stiffness is constrained by optimizing the objective function

$$\mathcal{L} = I - \lambda \sigma \quad (161)$$

for a given value of the Lagrange multiplier λ .

As shown in Fig. 5A, one-peaked solutions are more informative at low stiffness, while two-peaked solutions are more informative at high stiffness. We compute the convex hulls of the one- and two-peaked data to remove suboptimal solutions, and the transition occurs at the stiffness value at which the convex hulls intersect (cf. Fig. 5A). Repeating this procedure for many choices of gain allows one to trace out the phase transition shown Fig. 5B, which makes clear that one-peaked solutions are most informative at low stiffness, two-peaked solutions are most informative at high stiffness, and the critical stiffness decreases weakly with increasing gain.

IV. CONCLUSIONS

The presented spectral method exploits the linearity of the master equation to solve for probability distributions directly by expanding in the natural eigenfunctions the linear operator. We demonstrate the method on three models of gene expression: a single gene with multiple expression states, a gene regulatory cascade, and a model that combines multi-state expression with explicit regulation through binding of transcription factor proteins.

The spectral method permits huge computational gains over simulation. As demonstrated for all spectral expansions of the two-gene cascade (cf. Fig. 3), directly solving for the distribution via the spectral method is $\sim 10^7$ - 10^8 times faster than building the distribution from samples using a simulation technique. This massive speedup makes possible optimization and inference problems requiring full probability distributions that were not computationally feasible previously. For example, by optimizing information flow in a two-gene cascade in which both parent and child undergo two-state production, we reveal a transition from a one-peaked to a two-peaked joint probability distribution when constraints on protein number and timescale separation are relaxed. We emphasize that this optimization would not have been possible without the novel efficiency of the spectral method.

The spectral method also makes explicit the linear algebraic structure underlying the master equation. In many cases, such as in two-state bursting and the two-gene threshold regulation problem, this leads to analytic solutions. In general, such as shown in the case of the linear cascade, this leads to a set of natural bases for expansion of the generating function and reveals the features of each basis that are better suited to different types of problems. Specifically, bases in which the parent species is expanded in eigenfunctions are best when the parent distribution is Poissonian, and bases in which the parent is left in protein number space are best when the parent distribution is concentrated at low protein number. As well, bases in which the eigenfunctions of the child depend on the number of copies of the parent's protein are best suited for smooth regulation functions, whereas a basis in which the eigenfunctions of the child are pa-




FIG. 5: **A:** Mutual information I (cf. Eqn. 158) versus stiffness σ (cf. Eqn. 160) for fixed gain ($\gamma = 16$, cf. Eqn. 159), obtained by optimizing Eqn. 161 for λ values between 10^{-3} and 10^1 . Squares denote solutions whose joint distribution p_{nm} has one peak (cf. B, lower left inset), and dots denote solutions for which p_{nm} has two peaks (cf. B, upper right inset). Solid lines show the convex hulls of the one- and two-peaked solutions. Dotted lines indicate the stiffness value at which the hulls intersect and the stiffness values of the hull points to the left and right of the intersection. **B:** Phase diagram between one- and two-peaked optimal solutions in the gain-stiffness plane. Circles and left and right error bars at each gain are determined by the stiffness values at the intersection of the one- and two-peak convex hulls and at the hull points to the left and right of the intersection respectively (see dotted lines for the example case in A). Solid line shows a line of best fit. Insets show examples of one- (lower left) and two-peaked (upper right) optimal distributions p_{nm} .

parameterized by a constant is more numerically robust for sharp regulation functions such as thresholds. In all cases the linear algebraic structure of the spectral decomposition yields numerical prescriptions that greatly outperform simulation techniques. We anticipate that the computational speedup of the method, as well as the removal of the statistical obstacle of density estimation inherently

limiting simulation-based approaches, will make spectral methods such as those demonstrated here useful in addressing a wide variety of biological questions regarding accurate and efficient modeling of noisy information transmission in biological systems.

APPENDIX A

In this appendix we describe two ways to compute the mixed products $\langle n|j \rangle$ and $\langle j|n \rangle$ between the protein number states $|n\rangle$ and the eigenstates $|j\rangle$: by direct evaluation and by recursive updating.

The direct evaluation follows from Eqns. 9, 8, 20, and 10, and the fact that repeated derivatives of a product follow a binomial expansion. Introducing g as the rate parameter for the $|j\rangle$ states,

$$\langle n|j \rangle = \oint \frac{dx}{2\pi i} \langle n|x \rangle \langle x|j \rangle \quad (162)$$

$$= \oint \frac{dx}{2\pi i} \frac{e^{g(x-1)}(x-1)^j}{x^{n+1}} \quad (163)$$

$$= \frac{1}{n!} \partial_x^n \left[e^{g(x-1)}(x-1)^j \right]_{x=0} \quad (164)$$

$$= \frac{1}{n!} \sum_{\ell=0}^n \frac{n!}{\ell!(n-\ell)!} \partial_x^{n-\ell} \left[e^{g(x-1)} \right]_{x=0} \times \partial_x^\ell \left[(x-1)^j \right]_{x=0} \quad (165)$$

$$= \sum_{\ell=0}^n \frac{1}{\ell!(n-\ell)!} \left[g^{n-\ell} e^{-g} \right] \times \left[\frac{j!}{(j-\ell)!} (-1)^{j-\ell} \theta(j-\ell+1) \right] \quad (166)$$

$$= (-1)^j e^{-g} g^n j! \xi_{nj}, \quad (167)$$

where

$$\xi_{nj} = \sum_{\ell=0}^{\min(n,j)} \frac{1}{\ell!(n-\ell)!(j-\ell)!(-g)^\ell}. \quad (168)$$

Similarly, noting Eqns. 21 and 7,

$$\langle j|n \rangle = n!(-g)^j \xi_{nj}, \quad (169)$$

with ξ_{nj} as in Eqn. 168. Eqns. 169 and 167 clearly reduce to Eqns. 26-27 for the special case $j = 0$.

It is more computationally efficient to take advantage of the selection rules in Eqns. 12-15 and 22-25 to compute the mixed products recursively. For example, using Eqns. 22, 16, and 14,

$$\begin{aligned} \langle n|j+1 \rangle &= \langle n|\hat{b}^+|j \rangle = \langle n|(\hat{a}^+ - 1)|j \rangle \\ &= \langle n-1|j \rangle - \langle n|j \rangle, \end{aligned} \quad (170)$$

which can be initialized using $\langle n|0 \rangle = e^{-g} g^n / n!$ (cf. Eqn. 167) and updated recursively in j . Eqn. 170 makes clear that in n space the $(j+1)$ th mode is simply the (negative of the) discrete derivative of the j th mode.

Alternatively, Eqns. 15, 17, and 23 give

$$\begin{aligned} (n+1)\langle n+1|j\rangle &= \langle n|\hat{a}^-|j\rangle = \langle n|(\hat{b}^- + g)|j\rangle \\ &= j\langle n|j-1\rangle + g\langle n|j\rangle, \end{aligned} \quad (171)$$

which can be initialized using $\langle 0|j\rangle = (-1)^j e^{-g}$ (cf. Eqn. 167) and updated recursively in n .

One may similarly derive recursion relations for $\langle j|n\rangle$, i.e.

$$\langle j|n+1\rangle = \langle j-1|n\rangle + \langle j|n\rangle, \quad (172)$$

$$(j+1)\langle j+1|n\rangle = n\langle j|n-1\rangle - g\langle j|n\rangle, \quad (173)$$

initialized with $\langle j|0\rangle = (-g)^j/j!$ or $\langle 0|n\rangle = 1$ respectively (cf. Eqn. 169) and updated recursively in n or j respectively.

One may also use the full birth-death operator $\hat{b}^+\hat{b}^-$ to derive the recursion relations

$$(n+1)\langle n+1|j\rangle = (g+n-j)\langle n|j\rangle - g\langle n-1|j\rangle, \quad (174)$$

$$g\langle j|n+1\rangle = (g+n-j)\langle j|n\rangle - n\langle j|n-1\rangle, \quad (175)$$

initialized with $\langle 0|j\rangle = (-1)^j e^{-g}$ and $\langle 1|j\rangle = (-1)^j e^{-g}(g-j)$ (cf. Eqn. 167), and $\langle j|0\rangle = (-g)^j/j!$ and $\langle j|1\rangle = (-g)^j(1-j/g)/j!$ (cf. Eqn. 169), respectively, and updated recursively in n . We find Eqns. 174-175 are more numerically stable than Eqns. 170-173, as the former are two-term recursion relations while the latter are one-term recursion relations.

APPENDIX B

In the limit $g_- = 0$, Eqn. 54 reads

$$\begin{aligned} G(x) &= \frac{\omega_+}{\omega_+ + \omega_-} e^y \Phi[\omega_-, \omega_+ + \omega_- + 1; -y] \\ &\quad + \frac{\omega_-}{\omega_+ + \omega_-} \Phi[\omega_+, \omega_+ + \omega_- + 1; y], \end{aligned} \quad (176)$$

where $y = e^{g_+}(x-1)$. Using the fact that [47]

$$e^y \Phi[\alpha, \beta; -y] = \Phi[\beta - \alpha, \beta; y], \quad (177)$$

Eqn. 176 can be written

$$\begin{aligned} G(x) &= \frac{\omega_+}{\omega_+ + \omega_-} \Phi[\omega_+ + 1, \omega_+ + \omega_- + 1; y] \\ &\quad + \frac{\omega_-}{\omega_+ + \omega_-} \Phi[\omega_+, \omega_+ + \omega_- + 1; y], \end{aligned} \quad (178)$$

or, noting Eqn. 55 and the fact that $\Gamma(z+1) = z\Gamma(z)$ for any z ,

$$\begin{aligned} G(x) &= \sum_j \left(\frac{\omega_+}{\omega_+ + \omega_-} \frac{\Gamma(j + \omega_+ + 1)}{\Gamma(\omega_+ + 1)} \right. \\ &\quad \left. + \frac{\omega_-}{\omega_+ + \omega_-} \frac{\Gamma(j + \omega_+)}{\Gamma(\omega_+)} \right) \\ &\quad \times \frac{\Gamma(\omega_+ + \omega_- + 1)}{\Gamma(j + \omega_+ + \omega_- + 1)} \frac{y^j}{j!}, \end{aligned} \quad (179)$$

$$\begin{aligned} &= \sum_j \left(\frac{\omega_+}{\omega_+ + \omega_-} \frac{(j + \omega_+) \Gamma(j + \omega_+)}{\omega_+ \Gamma(\omega_+)} \right. \\ &\quad \left. + \frac{\omega_-}{\omega_+ + \omega_-} \frac{\Gamma(j + \omega_+)}{\Gamma(\omega_+)} \right) \\ &\quad \times \frac{(\omega_+ + \omega_-) \Gamma(\omega_+ + \omega_-)}{(j + \omega_+ + \omega_-) \Gamma(j + \omega_+ + \omega_-)} \frac{y^j}{j!}, \end{aligned} \quad (180)$$

$$= \sum_j \frac{\Gamma(j + \omega_+)}{\Gamma(\omega_+)} \frac{\Gamma(\omega_+ + \omega_-)}{\Gamma(j + \omega_+ + \omega_-)} \frac{y^j}{j!} \quad (181)$$

$$= \Phi[\omega_+, \omega_+ + \omega_-; y], \quad (182)$$

as in Eqn. 56.

The marginal p_n is given by

$$\langle n|G\rangle = \frac{1}{n!} \partial_x^n [G(x)]_{x=0} \quad (183)$$

(cf. Eqn. 10). Using Eqn. 56 and the derivative of the confluent hypergeometric function,

$$\partial_y^n \Phi[\alpha, \beta; y] = \frac{\Gamma(n + \alpha)}{\Gamma(\alpha)} \frac{\Gamma(\beta)}{\Gamma(n + \beta)} \Phi[\alpha + n, \beta + n; y], \quad (184)$$

one obtains Eqn. 57.

APPENDIX C

In this appendix, we fill in the details of the derivations of the equations of motion for the latter three of the four spectral bases discussed in Sec. II A.

The $|j, k_j\rangle$ basis

Projecting the conjugate state $\langle j, k_j|$ onto Eqn. 83 (in which dummy indices j and k are changed to j' and k' respectively) gives

$$\begin{aligned} 0 &= \sum_{j'k'} (j' + \rho k') \langle j|j'\rangle \langle k_j|k'_j\rangle G_{j'k'} \\ &\quad + \sum_{j'k'} \langle j|\hat{b}_n^+ \hat{\Gamma}_n|j'\rangle \langle k_j|k'_j\rangle G_{j'k'} \\ &\quad + \rho \sum_{j'k'} \langle k_j|\hat{b}_m^+|k'_j\rangle \langle j|\hat{\Delta}_n(j')|j'\rangle G_{j'k'}. \end{aligned} \quad (185)$$

From the orthonormality of states, the first term of Eqn. 185 simplifies to

$$\sum_{k'} (j + \rho k') \langle k_j | k'_j \rangle G_{jk'} = (j + \rho k) G_{jk}. \quad (186)$$

Recalling Eqn. 30, the product $\langle k_j | k'_j \rangle$ simplifies to

$$\langle k_j | k'_j \rangle = \frac{(-Q_{jj'})^{k-k'}}{(k-k')!} \theta(k-k'+1), \quad (187)$$

with $Q_{jj'} = \bar{q}_j - \bar{q}_{j'}$, whereupon Eqn. 185, separating the part of its second term which is diagonal in k from that which is subdiagonal and applying Eqn. 24 to its third term, becomes

$$\begin{aligned} 0 &= (j + \rho k) G_{jk} \\ &+ \sum_{j'} \Gamma_{j-1,j'} G_{j'k} \\ &+ \sum_{k' < k} \sum_{j'} \Gamma_{j-1,j'} \frac{(-Q_{jj'})^{k-k'}}{(k-k')!} G_{j'k'} \\ &+ \rho \sum_{k' < k} \sum_{j'} \Delta_{jj'} \frac{(-Q_{jj'})^{k-k'-1}}{(k-k'-1)!} G_{j'k'}, \end{aligned} \quad (188)$$

with $\Gamma_{jj'}$ as in Eqn. 76 and

$$\Delta_{jj'} = \langle j | \hat{\Delta}_n(j') | j' \rangle = \langle j | (\bar{q}_{j'} - \hat{q}_n) | j' \rangle \quad (189)$$

$$= \langle j | (\bar{q}_j - \hat{q}_n) | j' \rangle \quad (190)$$

$$= \sum_n \langle j | n \rangle (\bar{q}_j - q_n) \langle n | j' \rangle \quad (191)$$

(where the orthonormality of $|j\rangle$ states is used in going from Eqn. 189 to Eqn. 190). Defining $\ell = k - k'$ and

$$V_{jj'}^\ell = \frac{(-Q_{jj'})^\ell}{\ell!}, \quad (192)$$

Eqn. 188 can be written more compactly as Eqn. 87.

The $|n, k_n\rangle$ basis

Projecting the conjugate state $\langle n, k_n |$ onto Eqn. 93 (in which dummy indices n and k are changed to n' and k' respectively) gives

$$\begin{aligned} 0 &= \sum_{n'k'} (g_{n'} + n' + \rho k') \langle n | n' \rangle \langle k_n | k'_{n'} \rangle G_{n'k'} \\ &- \sum_{n'k'} g_{n'} \langle n | n' + 1 \rangle \langle k_n | k'_{n'} \rangle G_{n'k'} \\ &- \sum_{n'k'} n' \langle n | n' - 1 \rangle \langle k_n | k'_{n'} \rangle G_{n'k'} \\ &+ \rho \sum_{n'k'} \Delta_{n'} \langle n | n' \rangle \langle (k-1)_n | k'_{n'} \rangle G_{n'k'}, \end{aligned} \quad (193)$$

where $\Delta_n = \bar{q}_n - q_n$. Noting that, as in Eqn. 30,

$$\langle k_n | k'_{n\pm 1} \rangle = \frac{(-Q_n^\pm)^{k-k'}}{(k-k')!} \theta(k-k'+1), \quad (194)$$

where $Q_n^\pm = \bar{q}_n - \bar{q}_{n\pm 1}$, Eqn. 193 becomes

$$\begin{aligned} 0 &= (g_n + n + \rho k) G_{nk} \\ &- g_{n-1} \sum_{k' \leq k} \frac{(-Q_n^-)^{k-k'}}{(k-k')!} G_{n-1,k'} \\ &- (n+1) \sum_{k' \leq k} \frac{(-Q_n^+)^{k-k'}}{(k-k')!} G_{n+1,k'} \\ &+ \rho \Delta_n G_{n,k-1}. \end{aligned} \quad (195)$$

Separating the parts of the second and third term that are diagonal in k and defining $\ell = k - k'$ and

$$V_{n\ell}^\pm = \frac{(-Q_n^\pm)^\ell}{\ell!}, \quad (196)$$

Eqn. 195 becomes Eqn. 97.

The $|j, k_n\rangle$ basis

Substituting Eqn. 103 into Eqn. 58 at steady state gives

$$\begin{aligned} 0 &= \sum_{jk} G_{jk} \{ g_{n-1} \langle n-1, m | j, k_{n-1} \rangle \\ &+ (n+1) \langle n+1, m | j, k_{n+1} \rangle - (g_n + n) \langle n, m | j, k_n \rangle \\ &+ \rho [q_n \langle n, m-1 | j, k_n \rangle + (m+1) \langle n, m+1 | j, k_n \rangle \\ &- (q_n + m) \langle n, m | j, k_n \rangle] \}, \end{aligned} \quad (197)$$

or, in terms of raising and lowering operators (cf. Eqns. 14-15),

$$\begin{aligned} 0 &= \sum_{jk} G_{jk} \langle n, m | \{ \hat{a}_n^+ \hat{g}_n | j, k_{n-1} \rangle \\ &+ \hat{a}_n^- | j, k_{n+1} \rangle - (\hat{g}_n + \hat{a}_n^+ \hat{a}_n^-) | j, k_n \rangle \\ &+ \rho [\hat{a}_m^+ \hat{q}_n | j, k_n \rangle + \hat{a}_m^- | j, k_n \rangle \\ &- (\hat{q}_n + \hat{a}_m^+ \hat{a}_m^-) | j, k_n \rangle] \}. \end{aligned} \quad (198)$$

Using the definitions in Eqns. 64-68, Eqn. 198 can be written as Eqn. 105.

Using Eqns. 107-109, Eqn. 105 can be written

$$\begin{aligned} 0 &= -\langle n, m | \sum_{j'k'} (j' + \rho k') | j', k'_n \rangle G_{j'k'} \\ &- \langle n, m | \sum_{j'k'} \hat{b}_n^+ \hat{\Gamma}_n | j', k'_n \rangle G_{j'k'} \\ &- \langle m | \rho \sum_{j'k'} (\bar{q}_n - q_n) \langle n | j' \rangle \hat{b}_m^+ | k'_n \rangle G_{j'k'} \\ &+ \langle m | \sum_{j'k'} g_{n-1} \langle n-1 | j' \rangle | \delta_- k'_n \rangle G_{j'k'} \\ &+ \langle m | \sum_{j'k'} (n+1) \langle n+1 | j' \rangle | \delta_+ k'_n \rangle G_{j'k'}, \end{aligned} \quad (199)$$

where

$$|\delta_{\pm} k_n\rangle = |k_{n\pm 1}\rangle - |k_n\rangle, \quad (200)$$

and the dummy indices j and k have been changed to j' and k' respectively. Using Eqn. 194 to note that

$$\langle k_n | \delta_{\pm} k'_n \rangle = \frac{(-Q_n^{\pm})^{k-k'}}{(k-k')!} \theta(k-k'), \quad (201)$$

where $Q_n^{\pm} = \bar{q}_n - \bar{q}_{n\pm 1}$, we multiply Eqn. 199 by $\langle k_n | m \rangle$ and sum over m to obtain

$$\begin{aligned} 0 &= -\langle n | \sum_{j'} (j' + \rho k) | j' \rangle G_{j'k} \\ &\quad - \langle n | \sum_{j'} \hat{b}_n^+ \hat{\Gamma}_n | j' \rangle G_{j'k} \\ &\quad - \rho \sum_{j'} (\bar{q}_n - q_n) \langle n | j' \rangle G_{j',k-1} \\ &\quad + \sum_{j'} g_{n-1} \langle n-1 | j' \rangle \sum_{\ell=1}^k V_{n\ell}^- G_{j',k-\ell} \\ &\quad + \sum_{j'} (n+1) \langle n+1 | j' \rangle \sum_{\ell=1}^k V_{n\ell}^+ G_{j',k-\ell}, \end{aligned} \quad (202)$$

in which we exploit the completeness of $|m\rangle$ states, i.e. $\sum_m |m\rangle \langle m| = 1$, and $V_{n\ell}^{\pm}$ is as in Eqn. 99. Multiplying Eqn. 202 by $\langle j | n \rangle$, summing over n , and exploiting $\sum_n |n\rangle \langle n| = 1$ for the first two terms, we obtain Eqn. 110.

APPENDIX D

In this appendix we explicitly solve for G_{nk} in Eqn. 117 using the tridiagonal matrix, or Thomas [40], algorithm. We start by identifying the subdiagonal, diagonal, superdiagonal, and right hand side elements of Eqn. 117, respectively, as

$$A_n = g \quad (n = 1 \dots N), \quad (203)$$

$$B_n = -(\rho k + g + n) \quad (n = 0 \dots N), \quad (204)$$

$$C_n = n + 1 \quad (n = 0 \dots N-1), \quad (205)$$

$$R_n = -g\phi_k^- \delta_{nn_1} - n_1 \phi_k^+ \delta_{nn_0} \quad (n = 0 \dots N), \quad (206)$$

where N is the cutoff in protein count n and $n_1 = n_0 + 1$. Auxiliary variables are defined iteratively as

$$C'_0 = \frac{C_0}{B_0}, \quad (207)$$

$$C'_n = \frac{C_n}{B_n - C'_{n-1} A_n} \quad (n = 1 \dots N-1), \quad (208)$$

$$R'_0 = \frac{R_0}{B_0}, \quad (209)$$

$$R'_n = \frac{R_n - R'_{n-1} A_n}{B_n - C'_{n-1} A_n} \quad (n = 1 \dots N), \quad (210)$$

and the solution is obtained by backwards iteration with

$$G_N^k = R'_N, \quad (211)$$

$$G_{n-1}^k = R'_{n-1} - C'_{n-1} G_n^k \quad (n = N \dots 1) \quad (212)$$

(where k has been moved from subscript to superscript for ease of reading).

Computing the first few terms of Eqn. 208 reveals the pattern

$$C'_n = -(n+1) \frac{\eta_n^k}{\eta_{n+1}^k}, \quad (213)$$

where

$$\eta_n^k = \sum_{i=0}^n \frac{n!}{i!(n-i)!} g^{n-i} \prod_{\ell=0}^{i-1} (\rho k + \ell), \quad (214)$$

with the convention that $\prod_a^b[\cdot] = 1$ if $a > b$. Note that since $\prod_{\ell=0}^{i-1} (\rho k + \ell) = \Gamma(\rho k + i) / \Gamma(\rho k)$, we may also use the integral representation of the Gamma function to write

$$\eta_n^k = \frac{1}{\Gamma(\rho k)} \sum_{i=0}^n \frac{n!}{i!(n-i)!} g^{n-i} \int_0^t dt e^{-t} t^{(\rho k + i - 1)} \quad (215)$$

$$= \frac{1}{\Gamma(\rho k)} \int_0^t dt e^{-t} t^{(\rho k - 1)} \sum_{i=0}^n \frac{n!}{i!(n-i)!} g^{n-i} t^i \quad (216)$$

$$= \frac{1}{\Gamma(\rho k)} \int_0^t dt e^{-t} t^{(\rho k - 1)} (g + t)^n. \quad (217)$$

Using Eqn. 210 it is immediately clear that

$$R'_{n < n_0} = 0. \quad (218)$$

The first nonzero term is

$$R'_{n_0} = \frac{n_1 \phi_k^+}{g n_0} \frac{\eta_{n_0}^k}{\eta_{n_0-1}^k} \frac{1}{\epsilon_{n_0}^k - 1}, \quad (219)$$

where we have defined

$$\epsilon_n^k = \frac{\rho k + g + n}{g n} \frac{\eta_n^k}{\eta_{n-1}^k}. \quad (220)$$

Further iteration of Eqn. 210 makes clear that

$$R'_{n > n_0} = \frac{n_1}{g} f_k \prod_{i=n_0}^n \left(\frac{1}{i} \frac{\eta_i^k}{\eta_{i-1}^k} \frac{1}{\epsilon_i^k - 1} \right) \quad (221)$$

$$= \frac{n_1}{g} \frac{(n_0 - 1)!}{n!} \frac{\eta_n^k}{\eta_{n_0-1}^k} f_k \prod_{i=n_0}^n \frac{1}{\epsilon_i^k - 1}, \quad (222)$$

where

$$f_k = \phi_k^+ - \frac{g n_0}{n_1} \frac{\eta_{n_0-1}^k}{\eta_{n_0}^k} (\epsilon_{n_0}^k - 1) \phi_k^-. \quad (223)$$

Computing the first few terms of Eqn. 212 reveals that

$$G_{n>n_0}^k = (\epsilon_n^k - 1)R'_{n>n_0}F_n^k \quad (224)$$

$$= \frac{n_1}{g} \frac{(n_0 - 1)!}{n!} \frac{\eta_n^k}{\eta_{n_0-1}^k} f_k F_n^k \prod_{i=n_0}^{n-1} \frac{1}{\epsilon_i^k - 1}, \quad (225)$$

where

$$F_n^k = \sum_{i=0}^{N-n} \prod_{\ell=n}^{n+i} \frac{1}{\epsilon_\ell^k - 1}. \quad (226)$$

At the threshold Eqn. 212 gives

$$G_{n_0}^k = \frac{n_1}{g n_0} \frac{\eta_{n_0}^k}{\eta_{n_0-1}^k} \frac{1}{\epsilon_{n_0}^k - 1} (\phi_k^+ + f_k F_{n_1}^k) \quad (227)$$

and since $R'_{n<n_0} = 0$, the solution is easily completed using Eqn. 212, giving

$$G_{n<n_0} = G_{n_0} \prod_{i=1}^{n_0-n} (-C'_{n_0-i}) \quad (228)$$

$$= \frac{n_1}{g} \frac{(n_0 - 1)!}{n!} \frac{\eta_n^k}{\eta_{n_0-1}^k} \frac{\phi_k^+ + f_k F_{n_1}^k}{\epsilon_{n_0}^k - 1}. \quad (229)$$

These results are summarized in Eqns. 120-125.

Acknowledgments

A.M. was supported by National Science Foundation Grant DGE-0742450. A.M. and C.W. were supported by National Science Foundation Grant ECS-0332479. A.M.W. was supported by the Princeton Center for Theoretical Science Fellowship and by Columbia's Professional Schools Diversity Short-Term Visiting Fellowship. C.W. was supported by National Institutes of Health Grants 5PN2EY016586-03 and 1U54CA121852-01A1.

-
- [1] S. Hooshangi, S. Thiberge, and R. Weiss, *Proc Natl Acad Sci USA* **102**, 3581 (2005).
 - [2] M. Thattai and A. van Oudenaarden, *Biophys J* **82**, 2943 (2002).
 - [3] M. B. Elowitz, A. J. Levine, E. D. Siggia, and P. S. Swain, *Science* **297**, 1183 (2002).
 - [4] E. M. Ozbudak, M. Thattai, I. Kurtser, A. D. Grossman, and A. V. Oudenaarden, *Nat Genet* **31**, 69 (2002).
 - [5] P. S. Swain, M. B. Elowitz, and E. D. Siggia, *Proc Natl Acad Sci USA* **99**, 12795 (2002).
 - [6] J. Paulsson, O. G. Berg, and M. Ehrenberg, *Proc Natl Acad Sci USA* **97**, 7148 (2000).
 - [7] J. M. Pedraza and A. V. Oudenaarden, *Science* **307**, 1965 (2005).
 - [8] J. Paulsson, *Nature* **427**, 415 (2004).
 - [9] G. Tkačik, C. G. Callan, and W. Bialek, *Phys. Rev. E* **78**, 11910 (2008).
 - [10] G. Tkacik, C. G. Callan, and W. Bialek, *Proc Natl Acad Sci USA* **105**, 12265 (2008).
 - [11] E. Ziv, I. Nemenman, and C. H. Wiggins, *PLoS ONE* **2**, e1077 (2007).
 - [12] A. Mugler, E. Ziv, I. Nemenman, and C. H. Wiggins, *arXiv q-bio.MN* (2008), 0811.2834v1.
 - [13] E. Emberly, *Phys. Rev. E* **77**, 41903 (2008).
 - [14] F. Tostevin and P. R. ten Wolde, *arXiv q-bio.MN* (2009), 0901.0280v2.
 - [15] Y. Lan and G. A. Papoian, *J. Chem. Phys.* **125**, 154901 (2006).
 - [16] C. Dellago, P. G. Bolhuis, F. S. Csajka, and D. Chandler, *Journal of Chemical Physics* **108**, 1964 (1998).
 - [17] C. Dellago, P. Bolhuis, and P. Geissler, *Adv Chem Phys* **123**, 1 (2002).
 - [18] T. S. van Erp and P. G. Bolhuis, *Journal of Computational Physics* **205**, 157 (2005).
 - [19] T. S. van Erp, D. Moroni, and P. G. Bolhuis, *Journal of Chemical Physics* **118**, 7762 (2003).
 - [20] J. Elf, G.-W. Li, and X. S. Xie, *Science* **316**, 1191 (2007).
 - [21] P. J. Choi, L. Cai, K. Frieda, and X. S. Xie, *Science* **322**, 442 (2008).
 - [22] A. Raj, C. S. Peskin, D. Tranchina, D. Y. Vargas, and S. Tyagi, *Plos Biol* **4**, e309 (2006).
 - [23] I. Golding, J. Paulsson, S. M. Zawilski, and E. C. Cox, *Cell* **123**, 1025 (2005).
 - [24] S. Tănase-Nicola, P. B. Warren, and P. R. T. Wolde, *Phys Rev Lett* **97**, 68102 (2006).
 - [25] J. S. van Zon, M. J. Morelli, S. Tănase-Nicola, and P. R. T. Wolde, *Biophys J* **91**, 4350 (2006).
 - [26] J. S. V. Zon and P. R. T. Wolde, *J. Chem. Phys.* **123**, 234910 (2005).
 - [27] N. G. van Kampen, *Stochastic processes in physics and chemistry* (Amsterdam: North-Holland, 1992).
 - [28] A. M. Walczak, M. Sasai, and P. G. Wolynes, *Biophysical Journal* **88**, 828 (2005).
 - [29] Y. Lan, P. G. Wolynes, and G. A. Papoian, *J. Chem. Phys.* **125**, 124106 (2006).
 - [30] Y. Lan and G. A. Papoian, *Phys Rev Lett* **98**, 228301 (2007).
 - [31] A. B. Bortz, M. H. Kalos, and J. L. Lebowitz, *J Comput Phys* **17**, 10 (1975).
 - [32] D. T. Gillespie, *J Phys Chem* **81**, 2340 (1977).
 - [33] R. E. Bellman, *Adaptive Control Processes* (Princeton, NJ: Princeton University Press, 1961).
 - [34] A. M. Walczak, A. Mugler, and C. H. Wiggins, *Proceedings of the National Academy of Sciences* **106**, 6529

- (2009).
- [35] S. Iyer-Biswas, F. Hayot, and C. Jayaprakash, Phys. Rev. E **79**, 31911 (2009).
 - [36] M. Doi, Journal of Physics A: Mathematical and General **9**, 1465 (1976).
 - [37] Y. B. Zel'Dovich and A. A. Ovchinnikov, Soviet Journal of Experimental and Theoretical Physics **47**, 829 (1978).
 - [38] L. Peliti, Journal of Physics A: Mathematical and General **19**, L365 (1986).
 - [39] D. C. Mattis and M. L. Glasser, Reviews of Modern Physics **70**, 979 (1998).
 - [40] L. Thomas, Watson Sci. Comput. Lab. Rept., Columbia University, New York (1949).
 - [41] J. Lin, Information Theory, IEEE Transactions on **37**, 145 (1991).
 - [42] B. D. Gomperts, I. M. Kramer, and P. E. R. Tantham, *Signal transduction* (San Diego, CA: Academic Press, 2002).
 - [43] A. Y. Ting and D. Endy, Science **298**, 1189 (2002).
 - [44] P. B. Detwiler, S. Ramanathan, A. Sengupta, and B. I. Shraiman, Biophys J **79**, 2801 (2000).
 - [45] H. Bolouri and E. H. Davidson, Proc Natl Acad Sci USA **100**, 9371 (2003).
 - [46] B. L. Bassler, Curr Opin Microbiol **2**, 582 (1999).
 - [47] W. Koepf, *Hypergeometric summation: an algorithmic approach to summation and special function identities* (Braunschweig, Germany: Vieweg, 1998).
 - [48] A master equation in which the coordinate appears explicitly in the rates (e.g. g_n or q_n in Eqn. 58) is sometimes mistakenly termed “nonlinear” in the literature, perhaps discouraging calculations which exploit its inherent linear algebraic structure. We remind the reader that the master equation is perfectly linear.
 - [49] Note that setting $x = e^{ik}$ makes clear that the generating function is simply the Fourier transform.
 - [50] Note here, however, the difference with respect to the normalization convention commonly used in quantum mechanics in the prefactors of the creation and annihilation operations.
 - [51] Because this basis is a function of three coordinates (j , k , and n), we cannot abstractly define the generating function as in Eqn. 61; we must instead work directly with the joint probability distribution.